

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

152-17

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(31) International Patent Classification⁷:C07D 211/56, 417/14, 401/14, 413/14,
A61K 31/4468, 31/4523, A61P 31/12,
19/00

A1

(11) International Publication Number:

WO 00/65539

(43) International Publication Date:

9 November 2000 (09.11.00)

(21) International Application Number:

PCT/US00/11633

(22) International Filing Date:

1 May 2000 (01.05.00)

(30) Priority Date:

4 May 1999 (04.05.99)

US

(63) Reduced by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application

US
Filed on 09/205,187 (CIP)
4 May 1999 (04.05.99)(71) Applicant (*for all designated States except US*): SCHERING
CORPORATION [USUS]; 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).

(72) Inventor and

(73) Inventor/Applicants (*for US only*): BAROUDY, Bahije, M. [USUS]; 706 Central Avenue, Westfield, NJ 07090 (US). CLADER, John, W. [USUS]; 428 North Union Avenue, Cranford, NJ 07016 (US). JOSEPH, Hubert, B. [FRUS]; 5441 Washington Boulevard, Jersey City, NJ 07310 (US). MCCOMBS, Stuart, W. [GBUS]; 28 Hamilton Place, Caidwell, NJ 07006 (US). McKITTRICK, Brian, A. [USUS];


(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GR, HU, IL, IN, JP, KG, KR, KZ, LC, LK, LR, LT, LU, LV, MA, MD, ME, MK, MN, MX, NO, NZ, PL, PT, RO, RU, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, US, UZ, VN, YU, ZA, AR, IPO patent (OH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Euphrates patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BI, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(74) Agents: MAGATTI, Anita, W., et al.; Schering-Plough Corporation, Patent Department, K-6-1 1990, 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).

Published

With international search report.

(34) Title: PIPERIDINE DERIVATIVES USEFUL AS CCR5 ANTAGONISTS



5

PIPERIDINE DERIVATIVES USEFUL AS CCR5 ANTAGONISTS

BACKGROUND

The present invention relates to piperidine derivatives useful as selective CCR5 antagonists, pharmaceutical compositions containing the compounds, and methods of treatment using the compounds. The invention also relates to the use of a combination of a CCR5 antagonist of this invention and one or more antiviral or other agents useful in the treatment of Human Immunodeficiency Virus (HIV). The invention further relates to the use of a CCR-5 antagonist of this invention, alone or in combination with another agent, in the treatment of solid organ transplant rejection, graft v. host disease, arthritis, rheumatoid arthritis, inflammatory bowel disease, atopic dermatitis, psoriasis, asthma, allergies or multiple sclerosis.

The global health crisis caused by HIV, the causative agent of Acquired Immunodeficiency Syndrome (AIDS), is unquestioned, and while recent advances in drug therapies have been successful in slowing the progression of AIDS, there is still a need to find a safer, more efficient, less expensive way to control the virus.

It has been reported that the CCR5 gene plays a role in resistance to HIV infection. HIV infection begins by attachment of the virus to a target cell membrane through interaction with the cellular receptor CD4 and a secondary chemokine co-receptor molecule, and proceeds by replication and dissemination of infected cells through the blood and other tissue. There are various chemokine receptors, but for macrophage-tropic HIV, believed to be the key pathogenic strain that replicates *in vivo* in the early stages of infection, the principal chemokine receptor required for the entry of HIV into the cell is CCR5. Therefore, interfering with the interaction between the viral receptor CCR5 and HIV can block HIV entry into the cell. The present invention relates to small molecules which are CCR5 antagonists.

- 2 -


CCR-5 receptors have been reported to mediate cell transfer in inflammatory diseases such as arthritis, rheumatoid arthritis, atopic dermatitis, psoriasis, asthma and allergies, and inhibitors of such receptors are expected to be useful in the treatment of such diseases, and in the treatment of other inflammatory diseases or conditions such as inflammatory bowel disease, multiple sclerosis, solid organ transplant rejection and graft v. host disease.

Related piperidine derivatives which are muscarinic antagonists useful in the treatment of cognitive disorders such as Alzheimer's disease are disclosed in US patents 5,883,096; 6,037,352; 5,889,006; 5,952,349; and 5,977,138.


A-M. Vandamme et al., *Antiviral Chemistry & Chemotherapy*, 9:187-203 (1998) disclose current clinical treatments of HIV-1 infections in man including at least triple drug combinations or so-called Highly Active Antiretroviral Therapy ("HAART"). HAART involves various combinations of nucleoside reverse transcriptase inhibitors ("NRTI"), non-nucleoside reverse transcriptase inhibitors ("NNRTI") and HIV protease inhibitors ("PI"). In compliant drug-naïve patients, HAART is effective in reducing mortality and progression of HIV-1 to AIDS. However, these multidrug therapies do not eliminate HIV-1 and long-term treatment usually results in multidrug resistance. Development of new drug therapies to provide better HIV-1 treatment remains a priority.

SUMMARY OF THE INVENTION

The present invention relates to the treatment of HIV comprising administering to a human in need of such treatment an effective amount of a CCR5 antagonist represented by the structural formula I:




or a pharmaceutically acceptable salt thereof, wherein
X is -C(R¹³)₂-, -C(R¹³)(R¹⁹)-, -C(O)-, -O-, -NH-, -N(C₁-C₆alkyl)-,



- 5 -


unsubstituted carbon ring members, for example as shown in the following structures:



Preferred R⁷ and R⁸ substituents are: (C₁-C₆)alkyl, especially

5 methyl, halogen, especially chloro, and -NH₂. A preferred R⁹ substituent is hydrogen.

Also claimed are novel CCR5 antagonist compounds represented by the structural formula II



10 or a pharmaceutically acceptable salt thereof, wherein

(1) X^a is -C(R¹³)₂-, -C(R¹³)(R¹⁹)-, -C(O)-, -O-, -NH-, -N((C₁-C₆)alkyl)-,

OR³ CH₂-(C₁-C₆)alkyl-R³ NOR⁴ O-(C₁-C₆)alkyl CH₂-(C₁-C₆)alkyl
-CR¹³-, -CR¹³-, -C-, -CR¹³-, -C-

15 O-C(O)-(C₁-C₆)alkyl O-C(O)-O-(C₁-C₆)alkyl O-C(O)-NH-(C₁-C₆)alkyl
-CR¹³-, -CR¹³-, -CR¹³-

O-C(O)-N((C₁-C₆)alkyl)₂ NR⁵-C(O)-(C₁-C₆)alkyl
-CR¹³-

20 NR⁵-C(O)-O-(C₁-C₆)alkyl NR⁵-C(O)-NH-(C₁-C₆)alkyl
-CR¹³-

NR⁵-C(O)-N((C₁-C₆)alkyl)₂ C(O)-(C₁-C₆)alkyl
-CR¹³- or -N-

R^a is R^{6a}-phenyl, R^{6a}-pyridyl, R^{6a}-thiophenyl or R^{6a}-naphthyl;


R¹ is hydrogen, C₁-C₆ alkyl or C₂-C₆ alkenyl;

R² is R⁷, R⁸, R⁹-phenyl; R⁷, R⁸, R⁹-substituted 6-membered

25 heterocycle; R⁷, R⁸, R⁹-substituted 6-membered heterocycle N-oxide;

R¹⁰, R¹¹-substituted 5-membered heterocycle; naphthyl; fluorenyl;

- 6 -



R³ is R¹⁰-phenyl, pyridyl, pyrimidyl, pyrazinyl or thiazolyl;

R⁴ is hydrogen, C₁-C₆ alkyl, fluoro-C₁-C₆ alkyl, cyclopropylmethyl,

5 -CH₂CH₂OH, -CH₂CH₂-O-(C₁-C₆)alkyl, -CH₂C(O)-O-(C₁-C₆)alkyl,

-CH₂C(O)NH₂, -CH₂C(O)-NH(C₁-C₆)alkyl or -CH₂C(O)-N((C₁-C₆)alkyl)₂;

R⁵ and R¹¹ are independently selected from the group consisting of hydrogen and (C₁-C₆)-alkyl;

R^{6a} is 1 to 3 substituents independently selected from the group consisting of hydrogen, halogen, -CF₃, CF₃O-, -CN, -CF₃SO₂-, R²-phenyl,

10 -NHCOCF₃, 5-membered heterocycle and , wherein X is -O-, -NH- or -N(CH₃)-;

R⁶ is independently selected from the group consisting of R^{6a} and CH₃SO₂-;

15 (C₁-C₆)alkyl, halogen, -NR^aR^a, -OH, -CF₃, -OCH₃, -O-acyl, and -OCF₃;

R⁹ is R⁷, hydrogen, phenyl, -NO₂, -CN, -CH₂F, -CHF₂, -CHO,

-CH=NOR²⁰, pyridyl, pyrimidyl, N-oxide, pyrimidinyl, pyrazinyl,

-N(R²⁰)CONR²¹R²²-, -NHCONH(chloro-(C₁-C₆)alkyl), -NHCONH((C₃-C₁₀)-

cycloalkyl)(C₁-C₆)alkyl), -NHCO(C₁-C₆)alkyl, -NHCOCF₃, -NHCO₂N((C₁-

C₆)alkyl)₂, -NHCO₂(C₁-C₆)alkyl, -N(SO₂CF₃)₂, -NHCO₂(C₁-C₆)alkyl, C₃-C₁₀

cycloalkyl, -SR²³-, -SOR²³-, -SO₂R²³-, -SO₂NH(C₁-C₆)alkyl), -OSO₂(C₁-

C₆)alkyl, -OSO₂CF₃, hydroxy(C₁-C₆)alkyl, -CON R²⁰R²¹-, -CONCH₂CH₂-O-

CH₃)₂,

-OCONH(C₁-C₆)alkyl, -CO₂R²⁰-, -Si(CH₃)₃ or -B(OC(CH₃)₂)₂;

25 R¹⁰ is (C₁-C₆)alkyl, -NH₂ or R¹²-phenyl;

R¹² is 1 to 3 substituents independently selected from the group consisting of hydrogen, (C₁-C₆) alkyl, -CF₃, -CO₂R²⁰-, -CN, (C₁-C₆)alkoxy

and halogen;

R¹³, R¹⁴, R¹⁵ and R¹⁶ are independently selected from the group consisting of hydrogen and (C₁-C₆)alkyl;

30 consisting of hydrogen and (C₁-C₆)alkyl;

R¹⁷ and R¹⁸ are independently selected from the group consisting of hydrogen and C₁-C₆ alkyl, or R¹⁷ and R¹⁸ together are a C₂-C₅ alkylene group and with the carbon to which they are attached form a spiro ring of 3 to 6 carbon atoms;

- 7 -

R⁸ is R⁶-phenyl, R⁶-heteroaryl, R⁶-naphthyl, C₃-C₆ cycloalkyl, (C₃-C₆)cycloalkyl, (C₁-C₆)alkyl or (C₁-C₆)alkoxy(C₁-C₆)alkyl;

R²⁰, R²¹ and R²² are independently selected from the group consisting of H and C₁-C₆ alkyl; and

5 R²³ is C₁-C₆ alkyl or phenyl; or

(2):

X^a is -C(R¹³)(R¹⁹)-, -C(O)-, -O-, -NH-, -N((C₁-C₆)alkyl)-,

OR³ $\begin{array}{c} \text{CH}_2 \\ | \\ \text{C}_1\text{-C}_5\text{alkyl-R}^3 \\ | \\ \text{NOR}^{4a} \end{array}$ $\begin{array}{c} \text{O-C(O)-(C}_1\text{-C}_6\text{)alkyl} \\ | \\ \text{-CR}^{13}- \end{array}$ $\begin{array}{c} \text{O-C(O)-(C}_1\text{-C}_6\text{)alkyl} \\ | \\ \text{-CR}^{13}- \end{array}$

10 $\begin{array}{c} \text{O-C(O)-(C}_1\text{-C}_6\text{)alkyl} \\ | \\ \text{-CR}^{13}- \end{array}$ $\begin{array}{c} \text{O-C(O)-NH-(C}_1\text{-C}_6\text{)alkyl} \\ | \\ \text{-CR}^{13}- \end{array}$

$\begin{array}{c} \text{O-C(O)-N((C}_1\text{-C}_6\text{)alkyl)}_2 \\ | \\ \text{-CR}^{13}- \end{array}$ $\begin{array}{c} \text{NR}^5\text{-C(O)-(C}_1\text{-C}_6\text{)alkyl} \\ | \\ \text{-CR}^{13}- \end{array}$

15 $\begin{array}{c} \text{NR}^5\text{-C(O)-O-(C}_1\text{-C}_6\text{)alkyl} \\ | \\ \text{-CR}^{13}- \end{array}$ $\begin{array}{c} \text{NR}^5\text{-C(O)-NH-(C}_1\text{-C}_6\text{)alkyl} \\ | \\ \text{-CR}^{13}- \end{array}$

$\begin{array}{c} \text{NR}^5\text{-C(O)-N((C}_1\text{-C}_6\text{)alkyl)}_2 \\ | \\ \text{-CR}^{13}- \end{array}$ $\begin{array}{c} \text{C(O)-(C}_1\text{-C}_6\text{)alkyl} \\ | \\ \text{or -N-} \end{array}$

R^a is R^{6b}-phenyl, R^{6b}-pyridyl or R^{6b}-thiophenyl;

R^{4a} is fluoro-C₁-C₆ alkyl, cyclopropylmethyl, -CH₂CH₂OH,

20 -CH₂CH₂-O-(C₁-C₆)alkyl, -CH₂C(O)-O-(C₁-C₆)alkyl, -CH₂C(O)NH₂, -CH₂C(O)-NH-(C₁-C₆)alkyl or -CH₂C(O)-N((C₁-C₆)alkyl)₂;

R^{6b} is CH₃SO₂-; and

R¹, R², R³, R⁵, R¹⁴, R¹⁵ and R¹⁹ are as defined in (1).

25 Preferred are compounds of formula II(1) wherein R^a is R^{6a}-phenyl, especially wherein R^{6a} is a single substituent, and especially wherein the R^{6a} substituent is in the 4-position. Also preferred are compounds of formula II(1) wherein X^a is -CHOR³, -C(R³)(R⁹)- or -C(=NOR⁴)-; a preferred definition for R³ is pyridyl, especially 2-pyridyl, a preferred definition for R⁴ is (C₁-C₆)alkyl, especially methyl, ethyl or isopropyl, a preferred definition for R⁹ is hydrogen, and a preferred definition for R⁹ is R⁹-phenyl. For compounds of formula II(1), R¹ is preferably (C₁-C₆)alkyl,

- 8 -

especially methyl. Also for compounds of formula II(1), R¹⁴, R¹⁵ and R¹⁶ are preferably hydrogen.

Preferred are compounds of formula II(2) wherein R^a is R^{6b}-phenyl, especially wherein R^{6b} is a single substituent, and especially wherein the R^{6b} substituent is in the 4-position. Also preferred are compounds of formula II(2) wherein X^a is -CHOR³, -C(R³)(R⁹)- or -C(=NOR^{4a})-; a preferred definition for R³ is pyridyl, especially 2-pyridyl, preferred definitions for R^{4a} are cyclopropylmethyl and trifluoroethyl, a preferred definition for R⁹ is hydrogen, and a preferred definition for R⁹ is R⁹-phenyl.

10 For compounds of formula II(2), R¹ is preferably (C₁-C₆)alkyl, especially methyl. Also for compounds of formula II(2), R¹⁴, R¹⁵ and R¹⁶ are preferably hydrogen.

In compounds of formula II(1) and (2), R² is preferably R⁷, R⁸, R⁹-phenyl; R⁷, R⁸, R⁹-pyridyl or an N-oxide thereof, or R⁷, R⁸, R⁹-pyrimidyl. When R² is pyridyl, it is preferably 3- or 4-pyridyl, and when pyrimidyl, it is preferably 5-pyrimidyl. The R⁷ and R⁸ substituents are preferably attached to carbon ring members adjacent to the carbon joining the ring to the rest of the molecule and the R⁹ substituent can be attached to any of the remaining unsubstituted carbon ring members as shown above for compounds of formula I. Preferred R⁷ and R⁸ substituents for compounds of formula II are: (C₁-C₆)alkyl, especially methyl; halogen, especially chloro; and -NH₂; a preferred R⁹ substituent is hydrogen.

20 Another aspect of the invention is a pharmaceutical composition for treatment of HIV comprising an effective amount of a CCR5 antagonist of formula II in combination with a pharmaceutically acceptable carrier. Another aspect of the invention is a pharmaceutical composition for treatment of solid organ transplant rejection, graft v. host disease, arthritis, rheumatoid arthritis, inflammatory bowel disease, atopic dermatitis, psoriasis, asthma, allergies or multiple sclerosis comprising an effective amount of a CCR5 antagonist of formula II in combination with a pharmaceutically acceptable carrier.

30 Yet another aspect of this invention is a method of treatment of HIV comprising administering to a human in need of such treatment an effective amount of a CCR5 antagonist compound of formula II. Another aspect of the invention is a method of treatment of solid organ transplant rejection, graft v. host disease, arthritis, rheumatoid arthritis, inflammatory bowel disease, atopic dermatitis, psoriasis, asthma, allergies or multiple sclerosis

- 9 -

comprising administering to a human in need of such treatment an effective amount of a CCR5 antagonist compound of formula I or II.

- 5 Still another aspect of this invention is the use of a CCR5 antagonist of formula I or II of this invention in combination with one or more antiviral or other agents useful in the treatment of Human Immunodeficiency Virus for the treatment of AIDS. Still another aspect of this invention is the use of a CCR5 antagonist of formula I or II of this invention in combination with one or more other agents useful in the treatment of solid organ transplant rejection, graft v. host disease, inflammatory bowel disease, rheumatoid arthritis or multiple sclerosis. The CCR5 and antiviral or other agents which are components of the combination can be administered in a single dosage form or they can be administered separately; a kit comprising separate dosage forms of the actives is also contemplated.

15 DETAILED DESCRIPTION OF THE INVENTION

As used herein, the following terms are used as defined below unless otherwise indicated.

- Alkyl (including the alkyl portions of alkoxy, alkylamino and dialkylamino) represents straight and branched carbon chains and contains from one to six carbon atoms.

Alkenyl represents C₂-C₆ carbon chains having one or two unsaturated bonds, provided that two unsaturated bonds are not adjacent to each other.

- 25 Substituted phenyl means that the phenyl group can be substituted at any available position on the phenyl ring.

- AcyI means a radical of a carboxylic acid having the formula alkyl-C(O)-, aryl-C(O)-, aralkyl-C(O)-, (C₃-C₇)cycloalkyl-C(O)-, (C₃-C₇)cycloalkyl-(C₁-C₆)alkyl-C(O)-, and heteroaryl-C(O)-, wherein alkyl and heteroaryl are as defined herein; aryl is R¹²-phenyl or R¹²-naphthyl; and aralkyl is aryl-(C₁-C₆)alkyl, wherein aryl is as defined above.

- 30 Heteroaryl represents cyclic aromatic groups of 5 or 6 atoms or bicyclic groups of 11 to 12 atoms having 1 or 2 heteroatoms independently selected from O, S or N, said heteroatom(s) interrupting a carbocyclic ring structure and having a sufficient number of delocalized pi electrons to provide aromatic character, provided that the rings do not contain adjacent oxygen and/or sulfur atoms. For 6-membered heteroaryl rings, carbon atoms can be substituted by R⁷, R⁸ or R⁹ groups. Nitrogen atoms can form an N-oxide. All regioisomers are contemplated, e.g., 2-pyridyl, 3-

- 10 -

- pyridyl and 4-pyridyl. Typical 6-membered heteroaryl groups are pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl and the N-oxides thereof. For 5-membered heteroaryl rings, carbon atoms can be substituted by R¹⁰ or R¹¹ groups. Typical 5-membered heteroaryl rings are furyl, thienyl, pyrrolyl, thiazolyl, isothiazolyl, imidazolyl, pyrazolyl and isoxazolyl. 5-Membered rings having one heteroatom can be joined through the 2- or 3- position; 5-membered rings having two heteroatoms are preferably joined through the 4-position. Bicyclic groups typically are benzo-fused ring systems derived from the heteroaryl groups named above, e.g., quinolyl, phthalazinyl, quinoxalinyl, benzofuranyl, benzothienyl and indolyl.

- 10 Preferred points of substitution for 6-membered heteroaryl rings at R² are described above. When R² is a 5-membered heteroaryl group, the R¹⁰ and R¹¹ substituents are preferably attached to carbon ring members adjacent to the carbon joining the ring to the rest of the molecule, and R¹¹ is preferably alkyl; however, if a heteroatom is adjacent to the carbon joining the ring to the rest of the molecule (i.e., as in 2-pyrrolyl), R¹⁰ is preferably attached to a carbon ring member adjacent to the carbon joining the ring to the rest of the molecule.

Halogen represents fluoro, chloro, bromo and iodo.

- 20 Fluoro(C₁-C₆)alkyl represents a straight or branched alkyl chain substituted by 1 to 5 fluoro atoms, which can be attached to the same or different carbon atoms, e.g., -CH₂F, -CHF₂, -CF₃, F₃CCH₂- and -CF₂CF₃.

A therapeutically effective amount of a CCR5 antagonist is an amount sufficient to lower HIV-1-RNA plasma levels.

- 25 One or more, preferably one to four, antiviral agents useful in anti-HIV-1 therapy may be used in combination with a CCR5 antagonist of the present invention. The antiviral agent or agents may be combined with the CCR5 antagonist in a single dosage form, or the CCR5 antagonist and the antiviral agent or agents may be administered simultaneously or sequentially as separate dosage forms. The antiviral agents contemplated for use in combination with the compounds of the present invention comprise nucleoside and nucleotide reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors, protease inhibitors and other antiviral drugs listed below not falling within these classifications. In particular, the combinations known as HAART are contemplated for use in combination with the CCR5 antagonists of this invention.

35 The term "nucleoside and nucleotide reverse transcriptase inhibitors" ("NRTI"s) as used herein means nucleosides and nucleotides and

- 11 -

analogues thereof that inhibit the activity of HIV-1 reverse transcriptase, the enzyme which catalyzes the conversion of viral genomic HIV-1 RNA into proviral HIV-1 DNA.

- Typical suitable NRTIs include zidovudine (AZT) available under the RETROVIR tradename from Glaxo-Wellcome Inc., Research Triangle, NC 27709; didanosine (ddi) available under the VIDEX tradename from Bristol-Myers Squibb Co., Princeton, NJ 08543; zalcitabine (ddC) available under the HIVID tradename from Roche Pharmaceuticals, Nutley, NJ 07110; stavudine (d4T) available under the ZERT trademark from Bristol-Myers Squibb Co., Princeton, NJ 08543; lamivudine (3TC) available under the EPIVIR tradename from Glaxo-Wellcome Research Triangle, NC 27709; abacavir (1592U89) disclosed in WO96/30025 and available under the ZIAGEN trademark from Glaxo-Wellcome Research Triangle, NC 27709; didoxvir dipivoxil [bis(POM)-PMEA] available under the PREON tradename from Gilead Sciences, Foster City, CA 94404; lobucavir (BMS-180194), a nucleoside reverse transcriptase inhibitor disclosed in EP-0358154 and EP-0736533 and under development by Bristol-Myers Squibb, Princeton, NJ 08543; BCH-10652, a reverse transcriptase inhibitor (in the form of a racemic mixture of BCH-10618 and BCH-10619) under development by Biochem Pharma, Laval, Quebec H7V, 4A7, Canada; emilicilabine [(-)-FTC] licensed from Emory University under Emory Univ. U.S. Patent No. 5,814,639 and under development by Triangle Pharmaceuticals, Durham, NC 27707; beta-L-FD4 (also called beta-L-DAC and named beta-L-2', 3'-dideoxy-5-fluoro-cytidine) licensed by Yale University to Vion Pharmaceuticals, New Haven CT 06511; DAPD, the purine nucleoside, (-)-beta-D-2,6,-diamino-purine dioxolane disclosed in EP 0656778 and licensed by Emory University and the University of Georgia to Triangle Pharmaceuticals, Durham, NC 27707; and lodenosine (Fdda), 9-(2,3-dideoxy-2-fluoro-b-D-threo-pentofuranosyl)adenine, an acid stable purine-based reverse transcriptase inhibitor discovered by the NIH and under development by U.S. Bioscience Inc., West Conshohocken, PA 19428.

The term "non-nucleoside reverse transcriptase inhibitors" ("NNRTI"s) as used herein means non-nucleosides that inhibit the activity of HIV-1 reverse transcriptase.

Typical suitable NNRTIs include nevirapine (BI-RG-587) available under the VIRAMUNE tradename from Boehringer Ingelheim, the manufacturer for Roxane Laboratories, Columbus, OH 43216; delaviridine

- 12 -

- (BHAP, U-90152) available under the RESCRIPTOR tradename from Pharmacia & Upjohn Co., Bridgewater NJ 08807; efavirenz (DMP-266) a benzoxazin-2-one disclosed in WO94/03440 and available under the SUSTIVA tradename from DuPont Pharmaceutical Co., Wilmington, DE 19880-0723; PNU-142721, a tropyridine-thio-pyrimide under development by Pharmacia and Upjohn, Bridgewater NJ 08807; AG-1549 (formerly Shionogi # S-1153); 5-(3,5-dichlorophenyl)-thio-4-isopropyl-1-(4-pyridyl)methyl-H-imidazol-2-ylmethyl carbonate disclosed in WO 96/10019 and under clinical development by Agouron Pharmaceuticals, Inc., LaJolla CA 92037-1020; MKC-442 (1-(ethoxy-methyl)-5-(1-methylethyl)-6-(phenylmethyl)-2,4(1H,3H)-pyrimidinedione) discovered by Mitsubishi Chemical Co. and under development by Triangle Pharmaceuticals, Durham, NC 27707; and (+)-calanolide A (NSC-675451) and B, coumarin derivatives disclosed in NIH U.S. Patent No. 5,489,697, licensed to Med Chem Research, which is co-developing (+) calanolide A with Vita-Invest as an orally administrable product.

The term "protease inhibitor" ("PI") as used herein means inhibitors of the HIV-1 protease, an enzyme required for the proteolytic cleavage of viral polyprotein precursors (e.g., viral GAG and GAG Pol polyproteins), into the individual functional proteins found in infectious HIV-1. HIV protease inhibitors include compounds having a peptidomimetic structure, high molecular weight (7600 daltons) and substantial peptide character, e.g. CRIVIVAN (available from Merck) as well as nonpeptide protease inhibitors e.g., VIRACEPT (available from Agouron).

Typical suitable PIs include saquinavir (Ro 31-8959) available in hard gel capsules under the INVIRASE tradename and as soft gel capsules under the FORTOVASE tradename from Roche Pharmaceuticals, Nutley, NJ 07110-1199; ritonavir (ABT-538) available under the NORVIR tradename from Abbott Laboratories, Abbott Park, IL 60064; indinavir (MK-639) available under the CRIVIVAN tradename from Merck & Co., Inc., West Point, PA 19486-0004; nelfinavir (AG-1343) available under the VIRACEPT tradename from Agouron Pharmaceuticals, Inc., LaJolla CA 92037-1020; amprenavir (141W94), tradename AGENERASE, a non-peptide protease inhibitor under development by Vertex Pharmaceuticals, Inc., Cambridge, MA 02139-4211 and available from Glaxo-Wellcome, Research Triangle, NC under an expanded access program; lasinavir (BMS-234475) available from Bristol-Myers Squibb, Princeton, NJ 08543 (originally discovered by Novartis, Basel, Switzerland (CGP-61755); DMP-

- 13 -

- 450, a cyclic urea discovered by Dupont and under development by Triangle Pharmaceuticals; BMS-2322623, an azapeptide under development by Bristol-Myers Squibb, Princeton, NJ 08543, as a 2nd-generation HIV-1 PI; ABT-378 under development by Abbott, Abbott Park, IL 60064; and AG-1549 an orally active imidazole carbamate discovered by Shionogi (Shionogi #S-1153) and under development by Agouron Pharmaceuticals, Inc., LaJolla CA 92037-1020.

- Other antiviral agents include hydroxyurea, ribavirin, IL-2, IL-12, pentastuside and Yissum Project No. 11607. Hydroxyurea (Droxia), a ribonucleoside triphosphate reductase inhibitor, the enzyme involved in the activation of T-cells, was discovered at the NCI and is under development by Bristol-Myers Squibb. In preclinical studies, it was shown to have a synergistic effect on the activity of didanosine and has been studied with stavudine. IL-2 is disclosed in Ailomoto EP-0142268. Takeda EP. 0176299, and Chiron U. S. Patent Nos. RE 33653, 4530787, 4569790, 4604377, 4748234, 4752585, and 4949314, and is available under the PROLEUKIN (aldesleukin) tradename from Chiron Corp., Emeryville, CA 94608-2997 as a lyophilized powder for IV infusion or sc administration upon reconstitution and dilution with water; a dose of about 1 to about 20 million IU/day, sc is preferred; a dose of about 15 million IU/day, sc is more preferred. IL-12 is disclosed in WO96/25171 and is available from Roche Pharmaceuticals, Nutley, NJ 07110-1199 and American Home Products, Madison, NJ 07940; a dose of about 0.5 microgram/kg/day to about 10 microgram/kg/day, sc is preferred. Pentastuside (DP-178, T-20) a 36-amino acid synthetic peptide, disclosed in U.S. Patent No. 5,464,933 licensed from Duke University to Timeris which is developing pentastuside in collaboration with Duke University; pentastuside acts by inhibiting fusion of HIV-1 to target membranes. Pentastuside (3-100 mg /day) is given as a continuous sc infusion or injection together with efavirenz and 2 PIs to HIV-1 positive patients refractory to a triple combination therapy; use of 100 mg/day is preferred. Yissum Project No. 11607, a synthetic protein based on the HIV -1 Vif protein, is under preclinical development by Yissum Research Development Co., Jerusalem 91042, Israel. Ribavirin, 1- β -D-ribofuranosyl-1H-1,2,4-triazole-3-carboxamide, is available from ICN Pharmaceuticals, Inc., Costa Mesa, CA; its manufacture and formulation are described in U.S. Patent No. 4,211,771.

The term "anti-HIV-1 therapy" as used herein means any anti-HIV-1 drug found useful for treating HIV-1 infections in man alone, or as part of

- 14 -

- multidrug combination therapies, especially the HAART triple and quadruple combination therapies. Typical suitable known anti-HIV-1 therapies include, but are not limited to multidrug combination therapies such as (i) at least three anti-HIV-1 drugs selected from two NRTIs, one PI, a second PI, and one NNRTI; and (ii) at least two anti-HIV-1 drugs selected from NNRTIs and PIs. Typical suitable HAART - multidrug combination therapies include:

- (a) triple combination therapies such as two NRTIs and one PI; or (b) two NRTIs and one NNRTI; and (c) quadruple combination therapies such as two NRTIs, one PI and a second PI or one NNRTI. In treatment of naive patients, it is preferred to start anti-HIV-1 treatment with the triple combination therapy; the use of two NRTIs and one PI is preferred unless there is intolerance to PIs. Drug compliance is essential. The CD4⁺ and HIV-1-RNA plasma levels should be monitored every 3-6 months. Should viral load plateau, a fourth drug, e.g., one PI or one NNRTI could be added. See the table below wherein typical therapies are further described:

ANTI-HIV-1 MULTI DRUG COMBINATION THERAPIES

A. Triple Combination Therapies

1. Two NRTIs¹ + one PI²
2. Two NRTIs¹ + one NNRTI³

B. Quadruple Combination Therapies⁴

Two NRTIs + one PI + a second PI or one NNRTI

C. ALTERNATIVES⁵

- Two NRTI¹
- One NRTI⁵ + one PI²
- Two PIs⁶ ± one NRTI⁷ or NNRTI³
- One PI² + one NRTI⁷ + one NNRTI³

FOOTNOTES TO TABLE

1. One of the following: zidovudine + lamivudine; zidovudine + didanosine; stavudine + lamivudine; stavudine + didanosine; zidovudine + zalcitabine
2. Indinavir, nelfinavir, ritonavir or saquinavir soft gel capsules.
3. Nevirapine or delavirdine.
4. See A-M. Vandamme et al Antiviral Chemistry & Chemotherapy 9:187 at p 193-197 and Figures 1 + 2.

- 15 -


5. Alternative regimens are for patients unable to take a recommended regimen because of compliance problems or toxicity, and for those who fail or relapse on a recommended regimen. Double nucleoside combinations may lead to HIV-resistance and clinical failure in many patients.
6. Most data obtained with saquinavir and zidovudine (each 400 mg bid).
7. Zidovudine, stavudine or didanosine.
10. Agents known in the treatment of rheumatoid arthritis, transplant and graft v. host disease, inflammatory bowel disease and multiple sclerosis which can be administered in combination with the CCR5 antagonists of the present invention are as follows:
 - solid organ transplant rejection and graft v. host disease: immune suppressants such as cyclosporine and interleukin-10 (IL-10), tacrolimus, antilymphocyte globulin, OKT-3 antibody, and steroids;
 - inflammatory bowel disease: IL-10 (see US 5,368,854), steroids and azulfidine;
 - rheumatoid arthritis: methotrexate, azathioprine, cyclophosphamide, steroids and mycophenolate mofetil;
 - multiple sclerosis: interferon-beta, interferon-alpha, and steroids.
20. Certain CCR5 antagonist compounds of the invention may exist in different isomeric (e.g., enantiomers, diastereoisomers and atropisomers) forms. The invention contemplates all such isomers both in pure form and in admixture, including racemic mixtures.
25. Certain compounds will be acidic in nature, e.g., those compounds which possess a carboxyl or phenolic hydroxyl group. These compounds may form pharmaceutically acceptable salts. Examples of such salts may include sodium, potassium, calcium, aluminum, gold and silver salts. Also contemplated are salts formed with pharmaceutically acceptable amines such as ammonia, alkyl amines, hydroxyalkylamines, N-methylglucamine and the like.
30. Certain basic compounds also form pharmaceutically acceptable salts, e.g., acid addition salts. For example, the pyrido-nitrogen atoms may form salts with strong acid, while compounds having basic substituents such as amino groups also form salts with weaker acids. Examples of suitable acids for salt formation are hydrochloric, sulfuric, phosphoric,

- 16 -

- acetic, citric, oxalic, malonic, salicylic, malic, fumaric, succinic, ascorbic, maleic, methanesulfonic and other mineral and carboxylic acids well known to those in the art. The salts are prepared by contacting the free base form with a sufficient amount of the desired acid to produce a salt in the conventional manner. The free base forms may be regenerated by treating the salt with a suitable dilute aqueous base solution such as dilute aqueous NaOH, potassium carbonate, ammonia and sodium bicarbonate. The free base forms differ from their respective salt forms somewhat in certain physical properties, such as solubility in polar solvents, but the acid and base salts are otherwise equivalent to their respective free base forms for purposes of the invention.
- All such acid and base salts are intended to be pharmaceutically acceptable salts within the scope of the invention and all acid and base salts are considered equivalent to the free forms of the corresponding compounds for purposes of the invention.
- Compounds of the invention can be made by the procedures known in the art, for example by the procedures described in the following reaction schemes, by the methods described in the examples below, and by using the methods described in US patents 5,883,096; 6,037,352; 5,889,006; 5,952,349; and 5,977,138.
- The following solvents and reagents may be referred to herein by the abbreviations indicated: tetrahydrofuran (THF); ethanol (EtOH); methanol (MeOH); acetic acid (HOAc or AcOH); ethyl acetate (EtOAc); N,N-dimethylformamide (DMF); trifluoroacetic acid (TFA); trifluoroacetic anhydride (TFAA); 1-hydroxy-benzotriazole (HOBT); m-chloroperoxybenzoic acid (MCPBA); triethylamine (Et₃N); diethyl ether (Et₂O); tert-butoxycarbonyl (BOC); 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU); dimethylsulfoxide (DMSO); p-toluene sulfonic acid (p-TSA); potassium bis(trimethylsilyl)-amide (KHMDA); 4-dimethylaminopyridine (DMAP); N,N,N'-diisopropylethylamine (Dipea); and 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride (DEC). RT is room temperature.
- Compounds of formula I and II wherein X is $\text{CHO}(\text{C}=\text{O})-(\text{C}_1-\text{C}_6)\text{-alkyl}$, $\text{CHO}(\text{C}=\text{O})-(\text{C}_1-\text{C}_6)\text{-alkoxy}$, $\text{CHO}(\text{C}=\text{O})-\text{NH}-(\text{C}_1-\text{C}_6)\text{-alkyl}$, $\text{CHNR}^5(\text{C}=\text{O})-(\text{C}_1-\text{C}_6)\text{-alkyl}$, $\text{CHNR}^5(\text{C}=\text{O})-(\text{C}_1-\text{C}_6)\text{-alkoxy}$, $\text{CHNR}^5(\text{C}=\text{O})-\text{NH}-(\text{C}_1-\text{C}_6)\text{-alkyl}$ or $-\text{CHOR}^3$ (and wherein R¹⁴, R¹⁵ and R¹⁶ are hydrogen) are prepared according to Schemes 1-4:


- 17 -

Scheme 1




Compounds of formula 3, wherein R, R⁷ and R⁸ are as defined for formula 1, Z is CH or N, and R¹ is an alkyl group such as methyl were prepared as depicted in Scheme 1. Ketone 1, the synthesis of which was described in WO98/05292, was subjected to standard amidation with ArCOOH, EDCI or DEC, and HOBT, or ArCOCl, wherein Ar is R⁷, R⁸, substituted phenyl or pyridyl, followed by reduction with NaBH₄ to obtain 3. Derivatization of the free hydroxyl moiety with alkyl halides, acyl chlorides (R³COCl), alkyl chloroformates (ClCOOR³) and isocyanides (O=C=NR³) afforded ethers 4a, esters 4b, carbonates 4c, and carbamates 4d, respectively, wherein R³ is a lower alkyl group. The aryloxy compounds, 5, were obtained after condensation of the hydroxyl 3 with phenyl or pyridyl halides in the presence of a base.

Scheme 2




- 18 -



Alternatively, compounds of formula 5 can be prepared by reduction of the N-Boc ketone 1a to the alcohol 6 first, followed by functionalization of the free hydroxyl group with a halogen-substituted aryl in the presence of a base as shown in Scheme 2, or by a hydroxy-substituted aryl or heteroaryl (wherein Z¹ is as defined in Scheme 1) in the presence of PPh₃ and an azodicarboxylate of the formula R¹9O₂C=N=N-CO₂R²⁰, wherein R²⁰ is C₁-C₆ lower alkyl. Removal of the Boc protecting group and conversion to the amide is performed as in Scheme 1. This route allows the introduction of various aryloxy and heteroaryloxy moieties at R³ through the use of nucleophilic displacement or Mitsunobu-type reaction on intermediate 6.

Scheme 3




Compounds of formula 8, wherein R, R¹, R⁷, R⁸ and Z are as described in Scheme 1, were prepared by conversion of the ketone 2 to an oxime group with CH₃ONH₂·HCl, and reduction with BH₃·S(CH₃)₂ to

- 19 -

provide amine 8. Derivatization of the free amine moiety with an alkyl chloroformate (ClCOOR²⁰, wherein R²⁰ is C₁-C₆ alkyl) or an isocyanide (O=C=NR³) affords carbamate compounds 9 and urea compounds 10, respectively.


Scheme 4




Preparation of chiral analogs was performed through chemical resolution. The alcohol 6 was coupled with a chiral Boc-protected amino acid to obtain diastereoisomers 11a and 11b which were separated by chromatography. The chiral auxiliary was then removed with NaOH for each diastereoisomer and the same sequence of reactions described in Scheme 2 was carried out on each individual enantiomer to obtain compounds 12a and 12b.

Oximes of formula I or II wherein X is C=NO-R⁴ are prepared from the corresponding ketones from any of several methods known to those skilled in the art.

Scheme 5:




- 20 -



In Scheme 5, the ketone 1a, wherein R and R¹ are as defined for formula I and II, is dissolved in a solvent such as CH₃OH or ethanol and treated with an R⁴-substituted hydroxylamine such as O-methylhydroxylamine hydrochloride in the presence of a base such as sodium acetate.

The resulting mixture of Z- and E-O-substituted oximes 13 can be separated or the mixture carried through and separated at the end. The BOC protecting group is removed by treatment with an acid such as aqueous HCl or trifluoroacetic acid, and the resulting amine is coupled to an acid under standard conditions to obtain a compound of formula I or II.

Scheme 6:




Alternatively, the ketone 1a can be treated with HONH₂·HCl under similar conditions to yield, after separation, the E- and Z-oximes. Each oxime is then treated with a base such as potassium hexamethyldisilazide in a suitable solvent such as DMF followed by treatment with an alkylating agent, e.g., CH₃I, dimethylsulfate, CH₃CH₂I, trifluoroethyl triflate or similar electrophiles, to yield the desired O-substituted oxime.

The ketone starting material of formula 1a can be prepared by known methods as shown in Schemes 7 and 8.


- 21 -

Scheme 7:



In Scheme 7, Friedel-Crafts condensation of N-trifluoroacetyl-isonipicoloyl chloride 17 and an aromatic group R-H in the presence of a suitable catalyst such as AlCl_3 and optionally in a solvent such as CH_2Cl_2 yields a ketone 18 which is converted to its ethylene ketal 19 under standard conditions. The N-trifluoroacetyl group is removed and the resulting free amine 20 is treated with N-BOC-piperidine-4-one in the presence of a dehydrating agent such as titanium isopropoxide followed by treatment with diethylaluminum cyanide to give an aminonitrile 21. The aminonitrile is treated with a Grignard reagent ($\text{R}^1\text{Mg-halide}$) such as CH_3MgBr or vinylmagnesium bromide to give the alkylated product 22. The ketal is removed by treatment with aqueous acid followed by re-protection under standard conditions using BOC anhydride to give 1a.

Scheme 8:




Alternatively, 23, prepared via Wittig olefination of N-BOC-piperidone (Chen et al, *Tetrahedron Lett.*, 37, 30 (1996), 5233-5234), is transformed to

- 22 -

intermediate 25 by analogy to the procedure described in Scheme 7. 25 is converted to alcohol 26 by hydroboration/oxidation. Alcohol 26 is treated with a suitable oxidant such as a mixture of tetrapropylammonium perruthenate (TPAP) and N-methylmorpholine N-oxide (NMO) to give aldehyde 27. The aldehyde is treated with an aryllithium reagent in a suitable solvent such as ether or THF and the resulting alcohol 28 is treated with an oxidizing agent such as Dess-Martin periodinane or TPAP/NMO to give the desired ketone.


Compounds of formula I or II wherein X is -C(R¹³)(R¹⁹), wherein R and R¹⁹ are the same, or wherein R and R¹⁹ are different are prepared according to schemes 9 and 10, respectively. The schemes are exemplified by processes wherein R and R¹⁹ are each phenyl and wherein R is phenyl and R¹⁹ is CF₃-phenyl, respectively, but the general procedures apply to other R and R¹⁹ groups.

Scheme 9



N-BOC-4-piperidone is treated with CBr_4 to obtain the di-bromo compound of formula 44, which is then treated with phenylboronic acid to obtain the BOC-protected diphenylmethylene-piperidine of formula 45. The methylene bond is reduced using standard conditions to obtain the BOC-protected diphenylmethyl-piperidine of formula 46, the BOC group is removed and the amine of formula 47 is treated as described for compounds 20-22 of Scheme 7, the BOC group is removed by treatment with TFA, and the resultant amine subjected to a standard amidation procedure, e.g., treatment with a reagent R^2COOH and coupling agents such as EDCI, HOBt and a base, to obtain the compounds of formula 48.

Scheme 10




N-BOC-4-piperidone is treated with a reagent such as diethyl benzylphosphonate to obtain the phenylmethylene-piperidine of formula 49, which is then brominated to obtain the bromophenylmethylene-piperidine of formula 50. The BOC protecting group is removed using standard conditions, e.g., treatment with TFA, to obtain amine 51, and the amine 51 is treated as described for compounds 20-22 of Scheme 7 to obtain the aminonitrile 52, then the protected amine 53. The amine 53 is treated with a reagent such as 4-CF₃-phenylboronic acid to obtain compound 54 and the methylene bond is reduced using standard conditions to obtain racemic 55. The BOC group is removed by treatment with TFA, and the resultant amine subjected to a standard amidation procedure, e.g., treatment with a reagent R²COOH and coupling agents such as EDCI, HOBt and a base, to obtain the racemic compounds of formula 56.

Compounds useful in this invention are exemplified by the following preparative examples, which should not be construed to limit the scope of

the disclosure. Alternative mechanistic pathways and analogous structures within the scope of the invention may be apparent to those skilled in the art.

Example 1



A solution of free amine 29 (1.45 g, 3.97 mmol) and 2,6-dimethylbenzoyl chloride (840 mg, 5.0 mmol) in aqueous 1 N NaOH (20 ml) and CH₂Cl₂ (20 ml) was stirred overnight at RT. The reaction mixture was extracted with CH₂Cl₂, dried over Na₂SO₄ and concentrated under high vacuum to provide 30 (1.97 g, 97%), as a slightly yellow foam.

To a solution of ketone 30 (550 mg, 1.11 mmol) in CH₃OH (6 ml) was added NaBH₄ (60 mg, 1.59 mmol) and the solution was stirred overnight at RT. The reaction mixture was then poured into 0.1 N NaOH, extracted with CH₂Cl₂, dried over Na₂SO₄, and concentrated to give 31 (543 mg, 98%), as a slightly yellow foam.

Example 1A:

To a solution of alcohol 31 (50 mg, 0.10 mmol) in anhydrous DMF (0.5 ml) was added NaH (6.0 mg, 0.25 mmol) followed by ethyl iodide (12 μ l, 0.15 mmol) and the reaction was stirred 4 h at 40 °C. The reaction mixture was poured into aqueous 0.1 N NaOH, extracted with CH₂Cl₂, dried over Na₂SO₄, and concentrated. Purification by preparative chromatography (eluting with CH₂Cl₂/CH₃OH, 9:1) yielded 1A (31 mg, 59%) as a colorless oil: ¹H-NMR (300 MHz, CDCl₃) δ 7.39 (br d, J = 8.4 Hz, 2H), 7.02-7.12 (m, 3H), 6.95 (m, 2H), 3.94 (m, 1H), 3.79 (d, J = 7.2 Hz, 1H), 3.10-3.35 (m, 4H), 2.60-3.00 (m, 3H), 2.19 (br s, 6H), 1.60-2.10 (m, 5H), 1.05-1.50 (m, 5H), 1.08 (br t, 3H), 0.94 (s, 3H); HMMS (MH⁺) 527.2271.

- 25 -

Example 1B:

To a solution of alcohol **31** (50 mg, 0.10 mmol) and pyridine (16.2 μ l, 0.20 mmol) in anhydrous CH_2Cl_2 (0.5 mL) was added propionyl chloride (30 μ l, 0.30 mmol) and the solution was stirred overnight at RT. The reaction mixture was treated as for **1A** to give, after preparative chromatography (eluting with $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$, 9:1), **1B** (44.7 mg, 81%) as a colorless oil: ^1H -NMR (300 MHz, CDCl_3) δ 7.42 (br d, J = 8.2 Hz, 2H), 7.05-7.15 (m, 3H), 6.97 (m, 2H), 5.40 (d, J = 7.8 Hz, 1H), 4.09 (m, 1H), 3.43 (m, 1H), 3.23 (m, 1H), 2.96 (m, 1H), 2.82 (m, 1H), 2.70 (m, 1H), 2.21 (d, 3H), 1.60-2.10 (m, 5H), 1.05-1.45 (m, 5H), 1.08 (m, 3H), 0.95 (s, 3H); HRMS (MH^+) 555.2230.

Example 1C: To a solution of alcohol **31** (29.4 mg, 0.059 mmol) and pyridine (9.5 μ l, 0.118 mmol) in anhydrous CH_2Cl_2 (0.3 mL) was added methylchloroformate (13.8 μ l, 0.18 mmol) and the solution was stirred overnight at RT. The reaction mixture was treated as for **1A** to give, after preparative chromatography (eluting with $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$, 9:1), **1C** (15 mg, 46%) as a colorless oil: ^1H -NMR (300 MHz, CDCl_3) δ 7.46 (br d, J = 8.4 Hz, 2H), 7.14 (d, J = 8.4 Hz, 2H), 7.09 (m, 1H), 6.98 (m, 2H), 5.21 (d, J = 7.2 Hz, 1H), 4.09 (m, 1H), 3.71 (m, 3H), 3.45 (m, 1H), 3.24 (m, 1H), 2.97 (m, 1H), 2.82 (m, 1H), 2.70 (m, 1H), 2.22 (br s, 3H), 1.60-2.10 (m, 5H), 1.10-1.50 (m, 5H), 0.95 (s, 3H); HRMS (MH^+) 557.2017.

Example 1D:

A solution of alcohol **31** (30 mg, 0.060 mmol), pyridine (9.7 μ l, 0.12 mmol) and methylisocyanate (40 μ l, 0.68 mmol) in anhydrous THF (0.3 mL) was stirred 5 h at 45 $^\circ\text{C}$. The reaction mixture was treated as for **1A** to give, after preparative chromatography (eluting with $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$, 9:1), **1D** (25 mg, 75%) as a colorless oil: ^1H -NMR (300 MHz, CDCl_3) δ 7.42 (br d, J = 8.2 Hz, 2H), 7.05-7.15 (m, 3H), 6.98 (m, 2H), 5.34 (m, 1H), 4.08 (m, 1H), 3.44 (m, 1H), 3.24 (m, 1H), 3.19 (s, 3H), 2.96 (m, 1H), 2.65-2.85 (m, 2H), 2.20 (br s, 3H), 1.55-2.10 (m, 5H), 1.10-1.50 (m, 5H), 0.95 (s, 3H); HRMS (MH^+) 556.2169.


Example 1E:

A solution of alcohol **31** (50 mg, 0.10 mmol), NaH 60% in mineral oil (6 mg, 0.15 mmol), and 2-chloropyridine (28.2 μ l, 0.30 mmol) in anhydrous DMF (0.5 mL) was stirred 16 h at 90 $^\circ\text{C}$. The reaction mixture was treated as for **1A** to give, after preparative chromatography (eluting with $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$, 9:1), **1E** (50 mg, 86%) as a colorless oil: ^1H -NMR (300

- 26 -

MHz, CDCl_3) δ 7.98 (m, 1H), 7.47 (br t, J = 7.2 Hz, 1H), 7.38 (d, J = 8.0 Hz, 2H), 7.21 (d, J = 8.0 Hz, 2H), 6.95-7.15 (m, 3H), 6.65-6.80 (m, 2H), 5.74 (br d, J = 7.0 Hz, 1H), 4.09 (m, 1H), 3.44 (m, 1H), 3.24 (m, 1H), 2.65-3.05 (m, 3H), 2.22 and 2.23 (s, 3H), 1.60-2.15 (m, 5H), 1.10-1.50 (m, 5H), 0.87 (s, 3H); HRMS (MH^+) 576.2230.

Using similar procedures, compounds of the following structure were prepared



wherein R^3 , R^6 and R^2 are as defined in the table:

Ex.	R^6	R^3	R^2	HRMS (MH^+) found
1F	Br	-C(O)OCH ₂ CH ₃		571.2181
1G	Br	-C(O)CH ₃		541.2054
1H	Br	-C(O)-(CH ₂) ₂ CH ₃		569.2392
1I	Br	-C(O)NHCH ₂ CH ₃		572.2322
1J	Br			584.1786
1K	Br			577.2162
1L	Br			577.2183


Additional data for compounds of Example 1:

Ex.	^1H -NMR (300 MHz ^1H NMR (CDCl_3))
1J	7.49 (d, J = 8.4 Hz, 2H), 7.20-7.35 (m, 3H), 7.15 (m, 1H), 7.04 (m, 2H), 6.64 (d, J = 4.5 Hz, 1H), 5.58 and 5.60 (d, J = 7.2 Hz, 1H), 4.13 (m, 1H), 3.25-3.60 (m, 2H), 2.70-3.10 (m, 3H), 2.28 and 2.29 (s, 3H), 1.65-2.20 (m, 5H), 1.20-1.55 (m, 5H), 0.92 (br s, 3H)

- 27 -

1K	8.39 (d, $J = 5.6$ Hz, 1H), 7.42 (d, $J = 8.4$ Hz, 2H), 7.27 (d, $J = 8.4$ Hz, 2H), 7.05-7.20 (m, 2H), 6.99 (m, 2H), 6.84 (m, 1H), 5.70 (d, $J = 7.8$ Hz, 1H), 4.11 (m, 1H), 3.43 (m, 1H), 3.25 (m, 1H), 2.65-3.05 (m, 3H), 2.23 and 2.25 (s, 3H), 1.55-2.10 (m, 5H), 1.10-1.50 (m, 5H), 0.88 (br s, 3H)
----	--

Example 2



A solution of ketone **32** (0.60 g, 1.29 mmol) and NaBH_4 (60 mg, 1.59 mmol) in CH_3OH (5 ml) was stirred overnight at RT. The reaction mixture was poured into 0.1 N NaOH , extracted with CH_2Cl_2 , dried over Na_2SO_4 , and concentrated to give **33** (0.60 g, 100%), as a white foam.


To a solution of alcohol **33** (543 mg, 1.2 mmol) in anhydrous toluene (4 ml) was added KMnO_4 , 0.5 N in toluene (2.6 ml, 1.30 mmol) followed, 15 min. later, by 2-bromopyridine (125 μl , 1.30 mmol). The reaction was heated 5 h at 60 $^\circ\text{C}$, cooled to RT and poured into 5% aqueous NaHCO_3 (25 ml). Extraction with CH_2Cl_2 , drying over Na_2SO_4 and concentration afforded an oil which was purified by flash chromatography over silica gel (eluting with $\text{CH}_2\text{Cl}_2/\text{AcOEt}/\text{Et}_3\text{N}$ 50:50:1 to 40:60:1) to yield **34a** (310 mg, 49%), as a yellow foam.

A solution of **34a** (310 mg, 0.57 mmol) in anhydrous CH_2Cl_2 (2 ml) and TFA (2 ml) was stirred 30 min. at RT. After concentration, the residue was taken up in aqueous 1 N NaOH , extracted with CH_2Cl_2 , dried over Na_2SO_4 and concentrated to give **34b** (220 mg, 87%), as a white foam.

- 28 -

A solution of free amine **34b** (95 mg, 0.19 mmol), 2,4-dimethylnicotinic acid (50 mg, 1.45 mmol), DEC (60 mg, 0.31 mmol), HOBt (50 mg, 0.37 mmol) and N -methylmorpholine (80 ml, 0.72 mmol) in anhydrous DMF (1 ml) was stirred overnight at 40 $^\circ\text{C}$. After concentration, the residue was taken up in aqueous 0.1 N NaOH , extracted with CH_2Cl_2 , and dried over Na_2SO_4 . The residue obtained after concentration of the solvent was purified by preparative chromatography over silica gel (eluting with $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$, 96:4:1) to afford **35** (95 mg, 85%), as a colorless oil. $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 8.33 (d, $J = 5.1$ Hz, 1H), 7.99 (dd, $J = 4.8$ and 1.8 Hz, 1H), 7.86 (d, $J = 8.4$ Hz, 2H), 7.56 (d, $J = 8.4$ Hz, 2H), 7.53 (m, 1H), 6.96 (d, $J = 5.1$ Hz, 1H), 6.75-6.85 (m, 2H), 4.15 (m, 1H), 3.45 (m, 1H), 3.30 (m, 1H), 3.02 (s, 3H), 2.99 (m, 2H), 2.79 (m, 1H), 2.47 and 2.48 (s, 3H), 2.45 (m, 1H), 2.25 and 2.26 (s, 3H), 1.65-2.15 (m, 5H), 1.15-1.55 (m, 5H), 0.90 (s, 3H). HRMS (MH^+) 577.2658.

Using similar procedures, compounds of the following structure were prepared



wherein R^3 , R^6 and R^2 are as defined in the table:

Ex.	R^6	R^3	R^2	HRMS (MH^+) found
2A	Br			599.1062
2B	Br			578.2006
2C	Br			577.2172
2D	Br			577.2172
2E	H_3CSO_2			597.2296

- 29 -

2F	H ₃ CSO ₂ ⁻		578,2697
2G	F ₃ C-		567,2947
2H	H ₃ CSO ₂ ⁻		576,2890
2I	H ₃ CSO ₂ ⁻		593,2805
2J	F ₃ CO-		582,2969
2K	F ₃ CO-		584,2744
2L	F ₃ CO-		583,2913
2M	Br		580,2123
2N	Br		579,1986
2O	F ₃ CO-		599,2847
2P	Br		595,2114
2Q	Br		594,2072
2R	H ₃ CSO ₂ ⁻		578,2792

- 30 -

2S	H ₃ CSO ₂ ⁻		578,2801
2T	H ₃ CSO ₂ ⁻		594,2750
2U	H ₃ CSO ₂ ⁻		583,2426
2V	H ₃ CSO ₂ ⁻		576,2896
2W	H ₃ CSO ₂ ⁻		599,2362
2X	F ₃ C-		583,2805
2Y	F ₃ CO-		584,2848
2Z	F ₃ CO-		623,1790
2AA	Cl		533,2673
2BB	Cl		549,2646
2CC	Cl		573,1606
2DD	Cl		534,2637
2EE	Br		619,1062

- 31 -

2FF	H ₃ CSO ₂ -			584.2375
2GG	F ₃ C-			568.2913
2HH	H ₃ CSO ₂ -			618.1722
2II	H ₃ CSO ₂ -			579.2749
2JJ	F ₃ C-			607.1871
2KK	F			517.2696
2LL	F			533.2916
2MM	F			518.2944
2NN	Cl			589.1534
2OO	F			573.1818
2PP	Br			591.2330
2QQ	Br			607.2291

- 32 -

2RR	Br			592.2294
2SS	Br			633.1040
2TT	F ₃ C-			623.1809
2UU	F ₃ C-			583.2909
2VV	F ₃ C-			567.2961
2WW	F			532.3106
2XX	H			500.3023


Additional data for compounds of Example 2:

Ex.	¹ H-NMR (300 MHz ¹ H NMR (CDCl ₃))
2A	7.98 (m, 1H), 7.49 (br t, J = 7.1 Hz, 1H), 7.39 (d, J = 8.1 Hz, 2H), 7.12 (d, J = 8.1 Hz, 2H), 7.01 (t, J = 8.4 Hz, 1H), 6.65-6.80 (m, 3H), 6.56 (d, J = 8.4 Hz, 1H), 5.76 (d, J = 7.2 Hz, 1H), 3.95-4.20 (m, 1H), 3.89 and 3.92 (s, 2H), 3.30-3.55 (m, 2H), 3.12 (m, 1H), 2.70-3.00 (m, 2H), 1.65-2.10 (m, 5H), 1.20-1.60 (m, 5H), 0.95 and 0.99 (s, 3H)
2G	8.31 (d, 1H), 8.01 (d, 1H), 7.50 (m, 4H), 6.95 (d, 1H), 6.80 (m, 2H), 5.90 (d, 1H), 4.15 (d, 1H), 3.25-3.55 (m, 2H), 2.80-3.15 (m, 3H), 2.50 (d, 3H), 2.30 (d, 3H), 1.80-2.15 (m, 7H), 1.20-1.60 (m, 5H), 0.92 (s, 3H)

- 33 -



2K	7.97 (m, 1H), 7.45 (m, 1H), 7.32 (t, J = 8.4 Hz, 2H), 7.06 (m, 2H), 7.01 (m, 1H), 6.60-6.75 (m, 4H), 5.77 and 5.79 (d, J = 5.6 Hz, 1H), 3.55 (m, 1H), 3.32 (m, 1H), 2.70-2.95 (m, 2H), 2.18 (s, 3H), 1.65-2.10 (m, 5H), 1.15-1.55 (m, 5H), 0.78 and 0.91 (s, 3H)
2M	8.29 (d, J = 5.2 Hz, 1H), 8.18 (m, 1H), 7.98 (br s, 1H), 7.89 (br s, 1H), 7.38 (d, J = 8.4 Hz, 2H), 7.18 (d, J = 8.4 Hz, 2H), 6.92 (d, J = 5.2 Hz, 1H), 5.67 (d, J = 7.2 Hz, 1H), 4.07 (m, 1H), 3.43 (m, 1H), 3.26 (m, 1H), 2.65-3.05 (m, 3H), 2.41 and 2.42 (s, 3H), 2.20 (br s, 3H), 1.60-2.20 (m, 5H), 1.05-1.50 (m, 5H), 0.85 (br s, 3H)
2P	8.14 (d, J = 6.8 Hz, 1H), 8.02 (m, 1H), 7.51 (m, 1H), 7.41 (d, J = 8.0 Hz, 2H), 7.25 (d, J = 8.0 Hz, 2H), 6.98 (d, J = 6.4 Hz, 1H), 6.78 (m, 1H), 6.73 (m, 1H), 5.78 (d, J = 6.8 Hz, 1H), 4.17 (m, 1H), 3.43 (m, 1H), 3.32 (m, 1H), 2.95 (m, 1H), 2.86 (m, 1H), 2.75 (m, 1H), 2.44 and 2.46 (s, 3H), 2.23 and 2.25 (s, 3H), 1.65-2.10 (m, 5H), 1.15-1.50 (m, 5H), 0.90 (s, 3H)
2HH	8.49 (s, 2H), 8.26 (br s, 1H), 8.04 (br s, 1H), 7.80-7.95 (m, 3H), 7.53 (d, J = 8.4 Hz, 2H), 5.81 (d, J = 6.8 Hz, 1H), 4.16 (m, 1H), 3.30-3.50 (m, 2H), 2.94 (m, 2H), 2.80 (m, 1H), 1.75-2.15 (m, 5H), 1.25-1.50 (m, 5H), 0.89 (s, 3H)
2MM	8.93 (s, 1H), 8.04 (br d, J = 4.8 Hz, 1H), 7.50 (m, 1H), 7.32 (m, 2H), 6.97 (m, 2H), 6.78 (m, 1H), 6.72 (m, 1H), 5.82 (m, 1H), 4.21 (m, 1H), 3.25-3.50 (m, 2H), 2.93 (m, 2H), 2.78 (m, 1H), 2.44 and 2.46 (s, 3H), 1.90-2.15 (m, 3H), 1.70-1.90 (m, 2H), 1.15-1.50 (m, 5H), 0.90 (s, 3H)
2NN	8.17 (s, 1H), 8.01 (br d, J = 4.0 Hz, 1H), 7.50 (br t, J = 8.0 Hz, 1H), 7.20-7.35 (m, 4H), 6.78 (t, J = 6.8 Hz, 1H), 6.71 (m, 1H), 5.80 (d, J = 6.8 Hz, 1H), 4.18 (m, 1H), 3.44 (m, 1H), 3.39 (m, 1H), 3.00 (m, 2H), 2.80 (m, 1H), 1.70-2.15 (m, 5H), 1.10-1.50 (m, 5H), 0.90 (s, 3H)
2PP	8.37 (d, J = 6.0 Hz, 1H), 7.83 (br d, J = 4.6 Hz, 1H), 7.41 (d, J = 8.4 Hz, 2H), 7.34 (d, J = 6.0 Hz, 1H), 7.22 (d, J = 8.4 Hz, 2H), 6.97 (d, J = 4.6 Hz, 1H), 6.68 (br t, J = 6.0 Hz, 1H), 5.89 (br d, J = 6.8 Hz, 1H), 4.20 (m, 1H), 3.20-3.50 (m, 2H), 2.97 (m, 2H), 2.78 (m, 1H), 2.47 and 2.49 (s, 3H), 2.23 and 2.26 (s, 3H), 2.23 (s, 3H), 1.65-2.15 (m, 5H), 1.15-1.55 (m, 5H), 0.90 (s, 3H)

- 34 -

Example 3

- 35 -

3H), 3.47 (m, 1H), 3.25 (m, 1H), 2.88-3.10 (m, 3H), 2.25 (s, 6H), 1.20-2.10 (m, 12H), 0.90 (s, 3H); HRMS (MH⁺) 558.3013.

Example 4

5 A solution of alcohol **39ab** (660 mg, 1.41 mmol), Boc-Thr(t-Bu)-OH (413 mg, 1.50 mmol), DEC (290 mg, 1.50 mmol) and DMAP (190 mg, 1.55 mmol) in anhydrous CH₂Cl₂ (5 ml) was stirred overnight at RT. The reaction mixture was poured into aqueous saturated NaHCO₃, extracted with CH₂Cl₂, and dried over Na₂SO₄. The residue obtained after concentration of the solvent was subjected to flash chromatography over silica gel (eluting with CH₂Cl₂/acetone, 9:1) to afford, in order of elution: (i) first **40a** (391 mg, 38%), as a white foam; (ii) second **40b** (391 mg, 38%), as a white foam.

15 To a solution of diastereoisomer **40a** (391 mg, 0.54 mmol) in CH₃OH (3 ml) was added NaOH (110 mg, 2.75 mmol; 5 equiv.) and the solution was stirred at 65 °C for 3 h. The final mixture was then poured into aqueous 0.1 N NaOH and extracted with CH₂Cl₂ to yield **39a** (Enantiomer A) (246 mg, 98%) as a white foam. (Following the same procedure, **40b** gave **39b** (Enantiomer B). **40a** gives **43a** (Enantiomer A) and **40b** gives **43b** (Enantiomer B)).


- 36 -

5 A solution of alcohol **39a** (210 mg, 0.45 mmol), NaH 60% in mineral oil (23 mg, 0.96 mmol), and 2-bromopyridine (60 μ l, 0.62 mmol) in anhydrous DMF (1.5 ml) was stirred 2 h at 75 °C. The reaction mixture was poured into aqueous sat'd NaHCO₃, extracted with CH₂Cl₂, dried over Na₂SO₄ and purified by flash chromatography over silica gel (eluting with CH₂Cl₂/AcOEt/Et₃N, 60:40:0.5 to 40:60:0.5) to afford **41a** (143 mg, 59%).

Removal of the Boc-protecting group in **41a** (93 mg, 0.17 mmol) proceeded as for **34b** to provide **42a** (68 mg, 91%), as a white foam.

10 The amine **42a** (50 mg, 0.11 mmol) was coupled with 4,6-dimethyl-pyrimidine-5-carboxylic acid following the conditions described for the synthesis of **35** to yield **43a** (28 mg, 44%). ¹H-NMR (300 MHz, CDCl₃) δ 8.92 (s, 1H), 8.02 (m, 1H), 7.51 (m, 1H), 7.51 (br t, J = 8.4 Hz, 1H), 7.41 (d, J = 8.4 Hz, 2H), 7.24 (d, J = 8.4 Hz, 2H), 6.78 (m, 1H), 6.73 (m, 1H), 5.78 (m, 1H), 4.19 (m, 1H), 3.41 (m, 1H), 3.36 (m, 1H), 2.94 (m, 1H), 2.78 (m, 1H), 2.44 and 2.46 (s, 3H), 1.65-2.15 (m, 5H), 1.15-1.50 (m, 5H), 0.90 (s, 3H); HRMS (MH⁺) 578.2140.

The following compounds were prepared via similar methods:



wherein R³, R⁶ and R² are as defined in the table:

Ex.	Enantiomer	R ⁶	R ³	R ²	HRMS (MH ⁺) found
4A	A	Br			577.2172
4B	B	Br			577.2162
4C	B	Br			578.2119
4D	A	F ₃ CO-			584.2864

- 37 -


4E	B	F ₃ CO-			583,2862
4F	A	F ₃ CO-			583,2904
4G	A	F ₃ CO-			599,2857
4H	A	F ₃ CO-			598,2994
4I	B	F ₃ CO-			598,3000
4J	A	Cl			534,2639
4K	A	Cl			548,2784
4L	B	Cl			534,2644
4M	B	Cl			548,2784
4N	A	F ₃ CO-			599,2947
4O	B	F ₃ CO-			599,2947

- 38 -

Additional data for compounds of Example 4:

Ex.	¹ H-NMR (300 MHz ¹ H NMR (CDCl ₃))
4G	8.05 (m, 1H), 7.97 (s, 2H), 7.53 (t, J = 7.5 Hz, 1H), 7.41 (d, J = 8.4 Hz, 2H), 7.16 (d, J = 8.4 Hz, 2H), 6.81 (t, J = 6.4 Hz, 1H), 6.76 (m, 1H), 5.87 (m, 1H), 4.19 (m, 1H), 3.30-3.50 (m, 2H), 2.99 (m, 2H), 2.79 (m, 1H), 2.20 and 2.22 (s, 3H), 1.70-2.15 (m, 5H), 1.15-1.50 (m, 5H), 0.91 (s, 3H)
4I	8.03 (m, 1H), 7.53 (m, 1H), 7.39 (d, J = 8.4 Hz, 2H), 7.14 (d, J = 8.4 Hz, 2H), 6.79 (t, J = 6.8 Hz, 1H), 6.73 (m, 1H), 5.87 (m, 1H), 4.19 (m, 1H), 3.42 (m, 1H), 3.37 (m, 1H), 2.98 (m, 2H), 2.80 (m, 1H), 2.41 and 2.43 (s, 3H), 1.90-2.15 (m, 3H), 1.70-1.90 (m, 2H), 1.20-1.50 (m, 5H), 0.91 (s, 3H)

Example 5



- 1) T trifluoroacetic anhydride (TFAA) (300 ml) is added to isonicipic acid (96 g) at 0°C and the reaction mixture is heated at reflux for 4h. Excess TFAA is removed under vacuo, the reaction mixture is taken up in EtOAc, washed with water and concentrated to give 160 g of the amide. 50 g of this amide is treated with SOCl₂ (300 ml) and the reaction mixture heated at reflux overnight. Excess thionyl chloride is then removed under vacuo to give 54 g of the acid chloride.
- 2) AlCl₃ (11g) is added slowly to a solution of the product of step 1 (10 g) in bromobenzene (40 ml) at ambient temperature and the reaction mixture is heated at reflux for 4 h. It is then cooled and poured into a mixture of conc. HCl and ice, and the product is extracted with EtOAc. The organic layer is separated and washed with water, half saturated NaHCO₃ solution and concentrated to give 16.21 g of the desired ketone.
- 3) The product of step 2 (16.21 g) is dissolved in toluene (200 ml) containing ethylene glycol (25 ml) and p-toluenesulfonic acid (0.5 g). The reaction mixture is heated at reflux with azeotropic removal of water until no further water is collected. The reaction mixture is concentrated to give 17.4 g of the desired ketal.

- 39 -


- 4) The crude product of step 3 (17.4 g) is dissolved in CH₃OH (100ml) and to this is added water (25 ml) and K₂CO₃ (12 g) and the reaction mixture is stirred at ambient temperature overnight. The reaction mixture is diluted with water and extracted with EtOAc. The organic layer is separated, washed with water and brine, and concentrated to give 12.55 g of the desired amine.
- 5) To a stirred solution of the product of step 4 (7.2 g, 23 mmol) and N-BOC-piperidine-4-one (4.8 g, 24 mmol) in 1,2-dichloroethane (20 ml) is added titanium isopropoxide (6.7 ml, 32.3 mmol) and the mixture is stirred for 12 h at RT. The reaction mixture is concentrated and a 1.0 M solution of diethyl aluminium cyanide (35 ml) is added at RT and stirred for 3 h. The reaction mixture is then diluted with EtOAc, quenched with water (5 ml) and stirred for 2 h. The mixture is then filtered through celite and the resulting filtrate is concentrated and chromatographed with 30 % EtOAc/hexanes to afford 7.3 g (63%) of the desired cyanide.
- 6) To a stirred solution of the product of step 5 (7.3 g, 14.03 mmol) in THF (100 ml) is added a 3.0M solution CH₃MgBr in Et₂O (14.0 ml, 42 mmol) at RT and the mixture is stirred for 2 h. The reaction mixture is then quenched with saturated aqueous NH₄Cl and extracted with CH₂Cl₂. The extracts are concentrated to afford 7.0 g of desired methylated compound.
- 7) The crude ketal of step 6 is dissolved in EtOAc (100 ml) and 6 N HCl (40 ml) and conc. HCl (10 ml) is added and the mixture stirred at RT for 24 h. The reaction mixture is then neutralised with 20%NaOH and extracted with EtOAc, dried and concentrated to yield 5.0 g (98%) of amine.
- 8) To a stirred solution of the product of step 7 (5.0 g, 13.6 mmol) in Et₂O (200 ml) is added 10% NaOH (50 ml) and BOC₂O, and the mixture is stirred at RT overnight. The layers are separated and the organic layer is washed with brine, dried, concentrated and chromatographed with 20% EtOAc/hexanes to yield 5.1 g (79%) of the desired product.
- 9) To a stirred solution of the product of step 8 (1.5 g, 3.22 mmol) in CH₃OH (50 ml) is added sodium acetate (5.0 g, 47 mmol) and O-Methyl hydroxylamine hydrochloride and the mixture is stirred at RT for 24 h. The resulting mixture is then poured into aqueous NaOH and extracted with CH₂Cl₂. The combined extracts are dried, concentrated and chromatographed to yield 1.5 g (94%) of oxime as a mixture of E and Z isomers.
- 10) To a stirred solution of the product of step 9 (1.5 g, 3.0 mmol) in CH₂Cl₂ (10 ml) is added TFA (3 mL) and the mixture is stirred at RT for 2 h.

- 40 -

The reaction mixture is concentrated and poured into 10% NaOH and extracted with CH₂Cl₂. The combined extracts are dried concentrated to afford 1.2 g (100%) of amine.

- 11) To stirred solution of the product of step 10 (1.3 g, 3.2 mmol) in CH₂Cl₂ is added 2,6-dimethylbenzoic acid (0.74 g, 4.96 mmol), EDCI (0.94 g, 4.94 mmol), DIPEA (0.84 g, 6.58 mmol) and HOBT (0.66g, 4.94 mmol) and the mixture is stirred for 12 h at RT. The reaction mixture is quenched with NaHCO₃ and extracted with CH₂Cl₂. The combined extracts are dried and concentrated to yield 1.6 g of oxime as a mixture of E and Z isomers.
- 10) The isomers are separated by chromatography by eluting with CH₂Cl₂:Et₂O (4:1) to afford 0.77 g of E isomer and 0.49 g of Z isomer.
- E isomer: 300 MHz-¹H NMR (CDCl₃) δ 7.5 (d, 2H), 7.23 (m, 2H), 7.10 (m, 1H), 6.90 (d, 2H), 4.03 (m, 1H), 3.90 (s, 3H), 3.55 (m, 1H), 3.20 (m, 3H), 3.00 (m, 3H), 2.82 (m, 1H), 2.24 (s, 3H), 2.23 (s, 3H), 2.15 (m, 3H), 1.80-1.20 (m, 5H), 0.92 (s, 3H); MS FAB+ observed= 526.2070; estimated = 526.2069
- Z isomer: 300 MHz-¹H NMR (CDCl₃) δ 7.50 (d, 2H), 7.15-6.95 (m, 5H), 4.15 (m, 1H), 3.80 (s, 3H), 3.45 (s, 3), 3.25 (s, 3H), 3.00 (m, 2H), 2.24 (s, 3H), 2.25 (s, 3H), 2.10 (m, 2H), 1.80-1.50 (m, 7H), 0.92 (s, 3H); MS FAB+ observed= 526.2072; estimated = 526.2069.

The following compounds were prepared via similar methods:



wherein X, R⁰ and R² are as defined in the table:

Ex.	R ⁰	X	R ²	HRMS (MH ⁺) found
5A (mixture EZ)	Br			529.1017
5B (mixture EZ)	Br			549.1023
5C	Br			542.2210
5D	Br			549.1011

- 41 -

SE	Br		529.1128
SF	Br		530.1020
SG	Br		529.1017
SH	Br		542.1997
SI	Br		541.2178
SJ	Br		527.2787
SK	Br		543.1000
SL	Br		528.1971
SM	Br		541.2194
SN	Br		542.2132
SO	Br		583.1061
SP	Br		595.1895
SQ	Br		596.1831
SR	Br		541.2188

- 42 -

SS	Br		597.4911
ST	Br		569.0909
SU	Br		571.2270
SV	Br		556.2291
SW	Br		557.2119
SX	Br		557.2124
SY	Br		570.2454
SZ	Br		671.0058
SAA	Br		568.2286
SBB	Br		556.2286
SCC	Br		527.2015
SDD	Br		592.1000
SEE	Br		656.9869

- 43 -

5FF	Br	$\text{CH}_3\text{CH}_2\text{O}-\text{N}(\text{C})-\text{C}-$		686.9989
5GG	Br	$\text{CH}_3\text{CH}_2\text{O}-\text{N}(\text{C})-\text{C}-$		556.2290
5HH	$\text{F}_3\text{C}-$	$\text{CH}_3\text{CH}_2\text{O}-\text{N}(\text{C})-\text{C}-$		546.3056
5II	$\text{F}_3\text{C}-$	$\text{CH}_3\text{CH}_2\text{O}-\text{N}(\text{C})-\text{C}-$		531.2956
5JU	$\text{F}_3\text{C}-$	$\text{CH}_3\text{CH}_2\text{O}-\text{N}(\text{C})-\text{C}-$		547.2902
5KK	$\text{F}_3\text{C}-$	$\text{H}_3\text{CO}-\text{N}(\text{C})-\text{C}-$		517.2812
5LL	Br	$\text{H}_3\text{C}-\text{O}-\text{N}(\text{C})-\text{C}-$		555.2336
5MM	Br	$\text{H}_3\text{C}-\text{O}-\text{N}(\text{C})-\text{C}-$		567.2327
5NN	Br	$\text{CH}_3(\text{CH}_2)_2\text{O}-\text{N}(\text{C})-\text{C}-$		555.2341
5OO	Br	$\text{CH}_3\text{CH}_2\text{O}-\text{N}(\text{C})-\text{C}-$		610.2016
5PP	$\text{F}_3\text{CO}-$	$\text{CH}_3\text{CH}_2\text{O}-\text{N}(\text{C})-\text{C}-$		616.2746
5QQ	$\text{F}_3\text{C}-$	$\text{CH}_3\text{CH}_2\text{O}-\text{N}(\text{C})-\text{C}-$		600.2788

- 44 -


5RR	Br	$\text{CH}_3\text{CH}_2\text{O}-\text{N}(\text{C})-\text{C}-$		593.2131
5SS	Br	$\text{CH}_3\text{CH}_2\text{O}-\text{N}(\text{C})-\text{C}-$		590.1995
5TT	Br	$\text{CH}_3\text{CH}_2\text{O}-\text{N}(\text{C})-\text{C}-$		627.1729
5UU	Br	$\text{CH}_3\text{CH}_2\text{O}-\text{N}(\text{C})-\text{C}-$		556.218
5VV	Br	$\text{H}_3\text{CO}-\text{N}(\text{C})-\text{C}-$		542.2002
5WW	Br	$\text{CH}_3\text{CH}_2\text{O}-\text{N}(\text{C})-\text{C}-$		555.2336
5XX	Br	$\text{CH}_3\text{CH}_2\text{O}-\text{N}(\text{C})-\text{C}-$		655.287
5YY	Br	$\text{CH}_3\text{CH}_2\text{O}-\text{N}(\text{C})-\text{C}-$		566.2407
5ZZ	Br	$\text{H}_3\text{CO}-\text{N}(\text{C})-\text{C}-$		603.2349
5AB	Br	$\text{CH}_3\text{CH}_2\text{O}-\text{N}(\text{C})-\text{C}-$		617.2488
5AC	Br	$\text{CH}_3\text{CH}_2\text{O}-\text{N}(\text{C})-\text{C}-$		640.2868

- 45 -

Additional data for compounds of Example 5:

Ex.	¹ H-NMR (300 MHz ¹ H NMR (CDCl ₃))
5J	7.50 (d, 2H), 7.15-6.95 (m, 5H), 4.15 (m, 1H), 3.80 (s, 3H), 3.45 (s, 3H), 3.25 (s, 3H), 3.00 (m, 2H), 2.24 (s, 3H), 2.25 (s, 3H), 2.10 (m, 2H), 1.80-1.50 (m, 7H), 0.92 (s, 3H)
5L	8.95 (s, 1H), 7.53 (d, J = 8.4 Hz, 2H), 7.10 (d, J = 8.4 Hz, 2H), 4.24 (m, 1H), 3.81 (s, 3H), 3.98 (m, 2H), 2.75-3.00 (m, 3H), 2.48 (s, 3H), 2.45 (s, 3H), 1.99-2.20 (m, 4H), 1.73 (m, 3H), 1.20-1.62 (m, 4H), 0.94 (s, 3H)
5N	8.92 (s, 1H), 7.45 (d, J = 9.0 Hz, 2H), 7.10 (d, J = 8.7 Hz, 2H), 4.21 (m, 1H), 4.02 (d, J = 6.9 Hz, 2H), 3.98 (m, 2H), 2.75-2.92 (m, 3H), 2.46 (s, 3H), 2.41 (s, 3H), 1.90-2.20 (m, 4H), 1.73 (m, 3H), 1.27-1.62 (m, 4H), 1.15 (t, J = 8.1 Hz, 3H), 0.93 (s, 3H)

Example 6

A) Preparation of intermediate 27 (Scheme 8 (R¹ = CH₃)).

- 1) 23 (40.0 g, 0.203 mol) is vigorously stirred in EtOAc (200 ml) and concentrated aqueous HCl (80 ml) for 1.5 h. The solution is concentrated, diluted with Et₂O (300 ml) and H₂O (150 ml), the aqueous layer is separated and the organic layer is extracted once with H₂O (20 ml). Combined aqueous layers are concentrated and the residue is dried 24 h under high vacuum to provide 26.7 g (84%) of a white solid. To this hydrochloride and N-tert-butoxycarbonyl-4-piperidone (43.8 g, 0.22 mol) in anhydrous ClCH₂CH₂Cl (80 mL) with 4 Å molecular sieves, are successively added DBU (33.2 ml, 0.22 mol) and titanium(IV) isopropoxide (65.5 ml, 0.22 mol) at 0 °C, the reaction mixture is allowed to warm to RT and is stirred overnight at RT. The mixture is then cooled to 0 °C and diethylaluminum cyanide, 1 N in toluene (260 ml, 0.26 mol) is added with vigorous stirring. The reaction is allowed to warm to RT and stirred an additional 3 h, after which are added CH₂Cl₂ (300 ml), EtOAc (300 ml), and Celite (50 g). The reaction mixture is cooled to 0 °C, water (40 ml) is added slowly with vigorous stirring and, after an additional 5 min, stirring at RT, the excess of water is quenched with Na₂SO₄. The final mixture is then

- 46 -

filtered over Celite, evaporated and subjected to flash chromatography over silica gel (eluting with Hexanes/EtOAc, 8:2), to provide 50.3 g (83%) of 24 as a colorless oil which solidifies upon standing.


- 2) To a solution of 24 (27.7 g, 90.6 mmol) in anhydrous THF (200 mL) at 0 °C is slowly added CH₃MgBr 3 M in Et₂O (91 mL, 3 equiv.) with vigorous stirring. After the addition, the reaction is allowed to warm to RT and stirred 3 h. The reaction is then poured into aqueous saturated NH₄Cl, extracted with Et₂O (4 times), washed with brine, dried over Na₂SO₄, and concentrated to give 27.1 g (100%) of 25 as a colorless oil.
- 3) To a solution of 25 (11.6 g, 39.3 mmol) in anhydrous THF (50 ml) at 0 °C is slowly added BH₃·S(CH₃)₂ 2 N in THF (14 mL, 28 mmol) and the solution is stirred 2 days at RT. The final mixture is concentrated to ca. 50 ml and slowly poured into ice-cooled EtOH/THF 1:1 (50 ml). After 15 min. at 0 °C, 50 ml of a pH 7 buffer solution are added, followed slowly by 30% H₂O₂ aqueous solution (50 ml). The reaction mixture is stirred overnight at RT, diluted with 1 N NaOH and extracted with CH₂Cl₂. Combined organic layers are dried over Na₂SO₄, concentrated, then subjected to flash chromatography over silica gel (eluting with EtOAc/EtOH, 8:2) to yield 9.69 g (79%) of 26 as a colorless oil.
- 4) A solution of 26 (11.2 g, 35.8 mmol) and N-methylmorpholine N-oxide (4.67 g, 39.4 mmol) in anhydrous CH₂Cl₂ (100 ml) is stirred 1 h at RT, cooled to 0 °C, and TPAP (885 mg) is added portionwise. The reaction is allowed to warm to RT and stirred 1 h. Additional N-methylmorpholine N-oxide (1.30 g, 11 mmol) and TPAP (300 mg) are then added to drive the reaction to completion after 1 h. The reaction mixture is filtered over Celite, concentrated, then subjected to flash chromatography over silica gel (eluting with CH₂Cl₂/acetone, 8:2 to 7:3) to provide 5.91 g (53%) of 27 as a yellow oil.
- B) Preparation of title compounds of Example 6.
 - 1) A solution of 1-bromo-4-(trifluoromethoxy)-benzene (4.20 mL, 28.0 mmol) in anhydrous THF (100 mL) is cooled to -78 °C and n-BuLi 2.5 N in hexanes (11.2 mL, 28.0 mmol) is added via syringe. The reaction mixture is allowed to warm to -50 °C for 10 min, cooled to -78 °C, and a solution of aldehyde 27 (6.20 g, 20.0 mmol) in anhydrous THF (15 mL) is added dropwise. After stirring 30 min at -78 °C, then 30 min at -20 °C, the solution is poured into half-brine and extracted with CH₂Cl₂ (3 x 100 mL). Combined organic layers are dried over Na₂SO₄, and concentrated to give 8.85 g (94%) of an alcohol as a yellow oil.

- 2) To a solution of the product of step 1 (8.85 g, 39.3 mmol) in CH_2Cl_2 (100 ml) at 0 °C is added Dess-Martin periodinane (19.70 g, 2.5 equiv.) and the reaction mixture is stirred 2 h at RT. An additional 8.0 g of Dess-Martin periodinane is added and the reaction is stirred for an additional 4 h. The solution is poured into a 1:1 mixture of aqueous saturated NaHCO_3 and aqueous saturated $\text{Na}_2\text{S}_2\text{O}_3$ (200 ml), stirred 10 min, extracted with CH_2Cl_2 , and dried over Na_2SO_4 . The residue obtained after concentration of the solvents is purified by flash chromatography over silica gel (eluting with hexanes/EtOAc, 7:3) to yield 5.48 g (63%) of the ketone as a yellow oil.
- 3) A solution of the product of step 2 (2.85 g, 6.05 mmol), $\text{HONH}_2\cdot\text{HCl}$ (2.08 g, 30 mmol), and AcONa (2.46 g, 30 mmol) in EtOH (50 mL) is heated at reflux under N_2 for 4 h. After evaporation of the solvent, the residue is taken up in aqueous 0.1 N NaOH and extracted with CH_2Cl_2 .
- 15 The residue obtained after evaporation of the solvents is subjected to flash chromatography over silica gel, to afford first the E-hydroxime (eluting with $\text{CH}_2\text{Cl}_2/\text{EtOAc}$, 7:3; 0.84 g; 29%), then the Z-hydroxime (eluting with $\text{CH}_2\text{Cl}_2/\text{EtOAc}$ 1:1; 1.10 g; 37%), both products as white solids.
- 4) To a suspension of Z-hydroxime (0.89 g, 1.84 mmol) in anhydrous DMF (5 ml) is slowly added KHMDA 0.5 N in toluene (4.0 ml, 2.02 mmol) at 0 °C, leading to the appearance of a yellow solution. After 2 min. at this temperature, dimethylsulfate (350 μl , 3.7 mmol) is slowly added and the solution is allowed to warm to RT and stirred 1 h. The mixture is poured into aqueous 0.1 N NaOH, extracted with CH_2Cl_2 , and dried over Na_2SO_4 .
- 25 The residue obtained after concentration of the solvents is purified by flash chromatography over silica gel (eluting with hexanes/EtOAc, 75:25) to afford 0.55 g (62%) of the Z-methoxime as a slightly yellow oil.
- 5) A solution of Z-methoxime (0.59 g, 1.18 mmol) in anhydrous CH_2Cl_2 (6 ml) and TFA (3 ml) is stirred 1 h at RT. After concentration, the residue is taken up in aqueous 1 N NaOH, extracted with CH_2Cl_2 , dried over Na_2SO_4 , and concentrated to give 0.47 g (100%) of the free amine as a white foam.
- 30
- 6) A solution of the product of step 5 (470 mg, 1.18 mmol), 2,4-dimethylnicotinic acid (220 mg, 1.45 mmol), DEC (280 mg, 1.45 mmol), HOBt (243 mg, 1.80 mmol) and N-methylmorpholine (0.33 ml, 3.0 mmol) in anhydrous DMF is stirred 14 h. After concentration, the residue is taken up in aqueous 0.1 N NaOH, extracted with CH_2Cl_2 , and dried over Na_2SO_4 .
- 35 The residue obtained after concentration of the solvent is purified by flash

chromatography over silica gel (eluting with $\text{CH}_2\text{Cl}_2/\text{acetone}$, 7:3 to 1:1) to afford 640 mg (100%) of a colorless oil.















- 1H-NMR (400 MHz, CDCl_3) δ 8.35 (d, J = 7.8 Hz, 1H), 7.25 (AB system, 4H), 6.98 (d, J = 7.8 Hz, 1H), 4.22 (m, 1H), 3.82 (s, 3H), 3.43 (m, 1H), 3.33 (m, 1H), 2.99 (m, 2H), 2.85 (m, 1H), 2.49 (s, 3H, atropisomer a) and 2.51 (s, 3H, atropisomer b), 2.26 (s, 3H, atropisomer a) and 2/28 (s, 3H, atropisomer b), 1.95-2.21 (m, 3H), 1.20-1.90 (m, 7H), 0.92 (s, 3H). HRMS ($\text{M}+\text{H}^+$) 533.2747.






Following steps B-4, B-5, and B-6 using the E-oxime yields the corresponding E-methoxime product.



wherein R^4 , R^6 and R^2 are as defined in the table:

Ex.	R^6	R^4	R^2	HRMS ($\text{M}+\text{H}^+$) found
6A	Br			554.3000
6B	Br			555.2335
6C	Br			556.2175
6D	Br			571.2284
6E	Br			570.2331
6F	Br			569.1000
6G	$\text{F}_3\text{CO}-$			601.2628
6H	$\text{F}_3\text{CO}-$			617.2549

6I	F ₃ CO-	-CH ₃		534.2708
6J	F ₃ CO-		602.2465	
6K	F ₃ CO-		602.2579	
6L	F ₃ CO-		589.3013	
6M	Cl	CH ₃ CH ₂ -		513.2633
6N	Cl	CH ₃ -		483.2516
6O	F ₃ C-	CH ₃ -		533.2758
6P	Cl	CH ₃ CH ₂ -		497.2683
6Q	Cl	CH ₃ CH ₂ -		513.2642
6R	Cl	CH ₃ CH ₂ -		498.2633
6S	F ₃ C-	CH ₃ -		518.2749
6T	Cl	CH ₃ CH ₂ -		537.1603
6U	F ₃ C-	CH ₃ -		557.1680
6V	F ₃ C-	CH ₃ CH ₂ -		571.1838

6W	Cl	CH ₃ CH ₂ -		555.8401
6X	Cl	CH ₃ CH ₂ -		497.2682
6Y	F ₃ CO-	CH ₃ -		548.2853
6Z	F ₃ CO-	CH ₃ CH ₂ -		562.3017
6AA	F ₃ CO-	CH ₃ CH ₂ -		563.2939

Additional data for compounds of Example 6:

Ex.	¹ H-NMR (300 MHz, ¹ H NMR (CDCl ₃))
5F	8.31 (d, 1H), 7.51 (d, 2H), 7.10 (d, 2H), 6.95 (d, 2H), 4.20 (m, 2H), 3.40 (d, 2H), 3.30 (m, 2H), 3.35 (m, 3H), 2.80-3.05 (m, 5H), 2.45 (d, 3H), 2.25 (d, 3H), 1.25-2.20 (m, 10H), 0.50 (m, 2H), 0.22 (m, 2H), 0.90 (s, 3H)
6G	8.34 (d, J = 5.1 Hz, 1H), 7.24 (br s, 4H), 6.96 (d, J = 5.1 Hz, 1H), 4.33 (q, J = 8.6 Hz, 2H), 4.13 (m, 1H), 3.45 (m, 1H), 3.30 (m, 1H), 2.98 (m, 2H), 2.82 (m, 1H), 2.46 and 2.49 (s, 3H), 2.41 (m, 1H), 2.24 and 2.27 (s, 3H), 2.10 (m, 2H), 1.96 (m, 1H), 1.15-1.90 (m, 7H), 0.92 (s, 3H)
6I	8.92 (s, 1H), 7.23 (br s, 4H), 4.11 (m, 1H), 3.79 (s, 3H), 3.30-3.45 (m, 2H), 2.97 (m, 2H), 2.81 (m, 1H), 2.45 and 2.42 (s, 6H), 2.40 (m, 1H), 1.90-2.20 (m, 3H), 1.15-1.90 (m, 7H), 0.92 (s, 3H)

Example 7

Alternate synthesis of the compounds of Example 6.


- 5 1) The product of Example 6, step B-2 (566 mg, 1.20 mmol) is treated with H₃CONH₂·HCl using conditions similar to those shown in Example 6,

step B-3. The resulting crude mixture of Z- and E-methoximes is separated on a preparative silica gel TLC plate (eluting with hexanes/ EtOAc, 80:20) to afford, in order of elution, first the E-methoxime (175 mg; 29%), then the Z-methoxime (175 mg; 29%), both products as oils.

- 5 2) The Z-methoxime (75 mg; 0.15 mmol) of step 1 is deprotected following conditions similar to those shown in Example 6, step B-5 and the resulting free amine (46 mg) is directly subjected to amidation with 2,4-dimethylpicolinic acid using conditions similar to those shown in Example 6, step B-6 to yield 50 mg (82%) of a colorless oil.

10

The following compounds are prepared via similar procedures:



wherein R⁴, R⁶ and R² are as defined in the table:

Ex.	R ⁶	R ⁴	R ²	HRMS (M ⁺) found
7A	F ₃ CO-	CH ₃ -		532.2795
7B	F ₃ CO-	CH ₃ -		553.2192
7C	F ₃ CO-	CH ₃ -		533.2730
7D	F ₃ CO-	CH ₃ CH ₂ -		546.2940
7E	F ₃ C-	CH ₃ -		516.2833
7F	F ₃ CO-	CH ₃ -		534.2571
7G (E isomer)	F ₃ C-	CH ₃ -		537.2234

7H	F ₃ C-	CH ₃ -		537.2234
7I	F ₃ C-	CH ₃ -		537.2234
7J	F ₃ CO-	CH ₃ CH ₂ -		567.2362
7K	F ₃ C-	CH ₃ -		517.2812
7L	F ₃ C-	CH ₃ CH ₂ -		532.2787
7M	F ₃ CO-	CH ₃ CH ₂ -		547.2888
7N	F ₃ CO-			572.3093
7O	F ₃ CO-	CH ₃ CH ₂ -		548.2732
7P (E isomer)	F ₃ C-	CH ₃ -		517.2831
7Q	F ₃ CO-	CH ₃ -		549.2686
7R	F ₃ CO-	CH ₃ CH ₂ -		590.2854
7S	F ₃ C-	CH ₃ CH ₂ -		531.1002
7T	F ₃ C-	CH ₃ CH ₂ -		547.1348
7U (E isomer)	F ₃ CO-	CH ₃ -		532.2784
7V	F ₃ CO-			576.3049
7W	F ₃ CO-	CH ₃ CH ₂ -		563.2855

- 53 -


7X	F ₃ CO-		573.3052
7Y	F ₃ CO-		574.2889
7Z	F ₃ CO-		641.1537
7AA	F ₃ CO-		573.1638
7BB	F ₃ CO-		587.1821
7CC	F ₃ CO-		548.2861
7DD	F ₃ CO-		589.1610
7EE	F ₃ CO-		603.1748
7FF	F ₃ CO-		562.3030
7GG	F ₃ CO-		617.1918
7HH	F ₃ CO-		577.3019

Additional data for compounds of Example 7:

Ex.	¹ H-NMR (300 MHz ¹ H NMR (CDCl ₃))
7H	7.55 (d, 2H), 7.30 (d, 2H), 7.15 (t, 1H), 6.75 (d, 1H), 6.60 (d, 1H), 4.25 (m, 2H), 3.80 (s, 3H), 3.40 (m, 2H), 2.80-3.20 (m, 3H), 2.40 (m, 1H), 1.40-2.20 (m, 13H), 0.90 (s, 3H)

- 54 -

7K	8.31 (d, 1H), 7.61 (d, 2H), 7.31 (d, 2H), 6.95 (d, 2H), 4.30 (m, 2H), 3.80 (3, 2H), 3.20-3.50 (m, 2H), 2.75-3.05 (m, 3H), 2.45 (d, 3H), 2.25 (d, 3H), 1.45-2.20 (m, 11H), 0.92 (s, 3H)
7O	8.11 (d, J = 6.8 Hz, 1H), 7.25 (br s, 4H), 6.94 (d, J = 6.8 Hz, 1H), 4.16 (m, 1H), 3.75 (s, 3H), 3.20-3.45 (m, 2H), 2.85-3.00 (m, 3H), 2.41 (d, J = 11.6 Hz, 3H), 2.45 (m, 1H), 2.20 (d, J = 11.6 Hz, 3H), 1.85-2.20 (m, 3H), 1.15-1.85 (m, 7H), 0.88 (s, 3H)
7R	7.13-7.30 (m, 5H), 7.14 (m, 1H), 6.95 (m, 1H), 4.13 (m, 1H), 4.03 (q, J = 7.1 Hz, 2H), 3.15-3.50 (m, 2H), 2.86-3.10 (m, 2H), 2.80 (m, 1H), 2.39 (m, 1H), 2.15-2.30 (m, 6H), 1.85-2.15 (m, 3H), 1.10-1.85 (m, 7H), 1.28 (t, J = 7.1 Hz, 3H), 0.88 (br s, 3H)
7S	8.31 (d, 1H), 7.61 (d, 2H), 7.32 (d, 2H), 6.95 (d, 2H), 4.25 (m, 2H), 4.05 (q, 2H), 3.20-3.50 (m, 2H), 2.80-3.15 (m, 3H), 2.45 (d, 3H), 2.25 (d, 3H), 1.45-2.20 (m, 9H), 1.20 (t, 3H), 0.90 (s, 3H)

Example 8

- 1) To a stirred solution of the product of Example 5, step 8 (0.500 g, 1.07 mmol) in DMF (25 ml) is added sodium methylmercaptide (0.113 g, 1.62 mmol) and the mixture is heated to 70° C for 12 h. The reaction mixture is then cooled to RT, diluted with Et₂O, washed with brine, dried and concentrated to yield 0.437 g (97%) of sulfide.
- 2) A solution of the product of step 1 (1.00 g; 2.31 mmol), H₃CONH₂-HCl (3.80 g, 46.2 mmol), and AcONa (3.79 g, 46.2 mmol) in EtOH (30 ml) is heated at reflux under N₂ for 4 h. After evaporation of the solvent, the residue is taken up in aqueous 0.1 N NaOH and extracted with CH₂Cl₂. The residue obtained after evaporation of the solvents is subjected to flash chromatography over silica gel, to afford first the E-oxime (eluting Et₂O/CH₂Cl₂, 1:4; 0.45 g; 24%), then the Z-oxime (0.25 g, 15%).
- 3) To a solution of Z-oxime (0.250 g, 0.543 mmol) of step 2 in CH₃OH (5 ml) is at 0° C is added oxone (1.00 g, 1.627 mmol in 5 ml of CH₃OH) and the mixture is stirred at 0° C for 4 h. The reaction is then quenched with

- 55 -

10% NaOH, concentrated, poured into water (10 ml) and extracted with CH_2Cl_2 , dried and concentrated to yield 0.220 g (82%) of sulfone.


4) To a stirred solution of the product of step 3 (0.300 g, 0.608 mmol) in CH_2Cl_2 (5 ml) is added TFA (1 ml) and the mixture is stirred at RT for 2 h.

5 The reaction mixture is concentrated, poured into 10% NaOH and extracted with CH_2Cl_2 . The combined extracts are dried and concentrated to afford 0.240 g (100%) of amine.

5) To stirred solution of the product of step 4 (0.45 g, 0.114 mmol) in CH_2Cl_2 is added 2,6-dimethylnicotinic acid (0.26 g, 0.172 mmol), DEC (0.33 g, 0.172 mmol), N,N-diisopropylethylamine (DIPEA) (0.2 ml) and HOBT (0.24g, 0.172 mmol) and the mixture is stirred for 12 h at RT. The reaction mixture is quenched with NaHCO_3 , extracted with CH_2Cl_2 , dried, concentrated and purified by preparative chromatography (20% EtOH/EtOAc) to afford 0.046 g (76%) of Z-oxime amide.

15 300 MHz ^{-1}H NMR (CDCl_3) δ 8.32 (d, 1H), 7.95 (d, 2H), 7.40 (d, 2H), 6.95 (d, 1H), 4.20 (m, 1H), 3.82 (s, 3H), 3.30-3.45 (m, 3H), 3.10 (s, 3H), 2.80-3.00 (m, 3H), 2.50 (d, 2H), 2.25 (d, 2H), 1.30-2.20 (m, 12H), 0.92 (s, 3H).

The following compounds were prepared in a similar manner:




20 wherein X, R⁶ and R² are as defined in the table:

Ex.	R ⁶	X	R ²	HRMS (M ⁺) found
8A (mixture EZ)				526.2753
8B				547.2135
8C	Br			549.2133
8D				541.2849
8E				557.2798

- 56 -

8F				543.2641
8G				527.2692
8H	F ₃ C-			532.2895
8I				542.2796


Example 9



5 Dissolve the starting amine (2.0 g, 5.7 mmol) in CHCl_3 (57 ml, = Stock solution A ~ 0.1M). Add 430 μl of stock solution A (0.043 mmol) to a slurry of 0.25 g (~ 0.22 mmol) of resin bound carbodilimide (prepared by reacting Argopore-Cl resin with 1-(3-dimethylaminopropyl)3-ethyl carbodilimide in DMF at 100 C) in DMF (2 ml) in a polyethylene SPE cartridge. To this mixture add 0.12 ml of a 1M solution of 5-methyl-3-(2-chlorophenyl)isoxazole-4-carboxylic acid in DMF (0.12 mmol), HOBT (86 μl of a 0.5M solution in DMF) and DMAP (25 μl of a 0.05M solution in DMF). Shake this mixture for 14 h, filter and add 0.3 g of Amberlyst-15 resin (~ 1.5 mmol) to the filtrate. Shake for 1 to 2 h, filter and wash the resin twice with each of the following solvents: THF, CH_2Cl_2 and CH_3OH , then wash with THF and CH_2Cl_2 . Treat the resin with 2M NH_3 in CH_3OH (1 time for 30 min, and 1 time for 5 min.). Combine and concentrate the filtrates under reduced pressure to afford the title compound. LCMS found M⁺ = 570, 572 (calculated MW 571); TLC R_f = 0.45 ($\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$ (95/5/0.5)).

- 57 -


Using a similar procedure, the following compounds were prepared



wherein R² is as defined in the table:

Ex.	R ²	Data	TLC R _f values
9A		LCMS: MH ⁺ = 538.1 R _t = 6.27 min	0.58
9B		MS m/e = 475.2, 477.2 (Electrospray)	
9C		LCMS: MH ⁺ = 606	0.57
9D		LCMS: MH ⁺ = 507.1 R _t = 6.39 min	0.49
9E		LCMS: MH ⁺ = 497.1 R _t = 6.32 min	0.48

Example 10



5

Step 1: To a solution of alcohol **39ab** (406 mg; 0.87 mmol), 3-hydroxy-pyridine (95.1 mg; 1 mmol) and PPh₃ (262 mg; 1 mmol) in anhydrous THF (2 ml) at 0 °C was added diethylazodicarboxylate (160 ml; 1 mmol) and the mixture was allowed to warm to RT overnight. The reaction was poured into 5% aqueous NaHCO₃, extracted with CH₂Cl₂, and dried over Na₂SO₄.

10

After concentration of the solvents, the resulting oil was purified by flash chromatography over silica gel (eluting CH₂Cl₂/CH₃OH 97:3 to 95:5) to afford the desired compound (290 mg; 61%), as an oil.

Step 2: Removal of the Boc-protecting group of the product of step 1 (290 mg; 0.53 mmol) proceeded as in Example 2 to obtain the desired amine (210 mg; 89%), as a white foam.


15

- 58 -

Step 3: The amine of step 2 (50 mg; 0.11 mmol) was coupled with 4,6-dimethylpyrimidine-5-carboxylic acid following the conditions described in Example 2 to obtain the title compound (32 mg; 49%) as a colorless oil:
¹H-NMR (300 MHz, CDCl₃) δ 8.91 (s, 1H), 8.20 (br s, 1H), 8.10 (d, J = 4.5 Hz, 1H), 7.43 (br d, J = 8.4 Hz, 2H), 7.14 (br d, J = 8.4 Hz, 2H), 6.95-7.10 (m, 2H), 4.75 (br d, J = 6.8 Hz, 1H), 4.15 (m, 1H), 3.44 (m, 1H), 3.33 (m, 1H), 2.95 (m, 2H), 2.79 (m, 1H), 2.42 and 2.44 (s, 3H), 1.85-2.15 (m, 3H), 1.65-1.85 (m, 2H), 1.15-1.50 (m, 5H), 0.90 (s, 3H); HRMS (MH⁺) 578.2115.

10

Using similar procedures, compounds of the following structure were prepared



wherein R³, R⁶ and R² are as defined in the table:

Ex.	R ⁶	R ³	R ²	HRMS (MH ⁺) found
10A	CH ₃ SO ₂ -			592.2848
10B	Br			577.2166
10C	Br			595.2078
10D	F			517.2992
10E	F			516.3031
10F	F			532.2981
10G	Br			595.2072
10H	Cl			567.2308

- 59 -


10I	F ₃ C-		582.2955
10J	CH ₃ SO ₂ -		577.2853
10K	CH ₃ SO ₂ -		595.2764
10L	F ₃ CO-		601.2817
10M	F ₃ CO-		617.2514
10N	CH ₃ SO ₂ -		611.2460
10O	CH ₃ SO ₂ -		595.2749
10P	F ₃ C-		597.2951
10Q	F ₃ CO-		583.2905
10R	F ₃ CO-		598.2903
10S	F ₃ C-		601.2556
10T	F ₃ C-		585.2559
10U	F ₃ CO-		584.2860

- 60 -


Additional data for compounds of Example 10:

Ex.	¹ H-NMR (300 MHz ¹ H NMR (CDCl ₃))
10C	8.95 (s, 1H), 7.46 (br d, J = 8.4 Hz, 2H), 7.17 (br d, J = 8.4 Hz, 2H), 6.86 (t, J = 9 Hz, 2H), 6.70-6.72 (m, 2H), 4.69 (br d, J = 6.4 Hz, 1H), 4.19 (m, 1H), 3.47 (m, 1H), 3.37 (m, 1H), 2.99 (m, 2H), 2.82 (m, 1H), 2.47 and 2.50 (s, 3H), 1.90-2.15 (m, 3H), 1.65-1.90 (m, 2H), 1.20-1.50 (m, 5H), 0.93 (s, 3H)
10F	8.17 (d, J = 6.8 Hz, 1H), 7.28 (m, 2H), 7.18 (t, J = 7.5 Hz, 1H), 6.95-7.10 (m, 3H), 6.87 (t, J = 7.5 Hz, 1H), 6.80 (d, J = 7.5 Hz, 2H), 4.80 (d, J = 6.8 Hz, 1H), 4.17 (m, 1H), 3.25-3.50 (m, 2H), 2.99 (m, 2H), 2.80 (m, 1H), 2.43 (br s, 3H), 2.24 (br s, 3H), 1.65-2.20 (m, 5H), 1.15-1.50 (m, 5H), 0.90 (s, 3H)
10H	8.95 (s, 1H), 7.32 (br d, J = 8.4 Hz, 2H), 7.23 (br d, J = 8.4 Hz, 2H), 7.08 (t, J = 8.1 Hz, 1H), 6.80-6.90 (m, 2H), 6.68 (m, 1H), 4.77 (br d, J = 6.8 Hz, 1H), 4.19 (m, 1H), 3.46 (m, 1H), 3.37 (m, 1H), 3.00 (m, 2H), 2.81 (m, 1H), 2.47 and 2.49 (s, 3H), 1.90-2.15 (m, 3H), 1.65-1.90 (m, 2H), 1.20-1.50 (m, 5H), 0.93 (s, 3H)
10K	8.81 (s, 1H), 7.78 (d, J = 8.4 Hz, 2H), 7.53 (m, 1H), 7.47 (d, J = 8.4 Hz, 2H), 6.90 (m, 1H), 6.74 (m, 1H), 6.59 (m, 1H), 4.83 (d, J = 6.8 Hz, 1H), 4.08 (m, 1H), 3.20-3.40 (m, 2H), 2.70-3.00 (m, 3H), 2.35 (br s, 3H), 1.65-2.15 (m, 5H), 1.15-1.50 (m, 5H), 0.87 (s, 3H)
10L	8.33 (d, J = 5.1 Hz, 1H), 7.99 (dd, J = 4.8 and 1.8 Hz, 1H), 7.86 (d, J = 8.4 Hz, 2H), 7.56 (d, J = 8.4 Hz, 2H), 7.53 (m, 1H), 6.96 (d, J = 6.4 Hz, 1H), 6.75-6.85 (m, 2H), 4.15 (m, 1H), 3.45 (m, 1H), 3.30 (m, 1H), 3.02 (s, 3H), 2.99 (m, 2H), 2.79 (m, 1H), 2.47 and 2.48 (s, 3H), 2.45 (m, 1H), 2.25 and 2.26 (s, 3H), 1.65-2.15 (m, 5H), 1.15-1.55 (m, 5H), 0.90 (s, 3H)

Example 11



- 61 -



- 1) N-Boc-4-piperidone (10 g, 50 mmol) and PPh₃ (53 g, 200 mmol) were taken up in CH₃CN (100 ml). The solution was cooled to 0 °C and CBr₄ (33 g, 100 mmol) was added to the solution at 0 °C. The solution was stirred at 0 °C for 15 min. and at 25 °C for 2 h. Et₃O (200 ml) was added, and the resulting mixture was filtered through a plug of SiO₂.

Concentration gave a yellow solid. Purification via flash chromatography (9/1 hexanes/Et₂O, SiO₂) gave 10 g (56 %) of the di-bromo product as a white solid.

- 2) A solution of the product of step 1 (1 g, 2.8 mmol), PhB(OH)₂ (1.2 g, 9.9 mmol), PdCl₂(PPh₃)₂ (197 mg, 0.28 mmol), and Na₂CO₃ (897 mg, 8.5 mmol) were taken up in THF/H₂O (4/1, 20 ml) and stirred at 65 °C under N₂ for 24 h. The solution was partitioned between EtOAc and H₂O, the aqueous layer was extracted with EtOAc and the combined organic layers were washed with brine and dried over Na₂SO₄. Filtration and concentration gave a dark brown oil. Purification via flash chromatography (9/1 hexanes/Et₂O, SiO₂) gave 941 mg (96 %) of the desired product as a white solid, m.p. = 152-153 °C.

- 3) A solution of the product of step 2 (500 mg, 1.4 mmol) and Pd(OH)₂ on carbon (100 mg, 20 wt % Pd (dry basis), 50 wt % H₂O) were taken up in CH₃OH (20 ml) and shaken in a Parr apparatus under H₂ (50 psi) for 15 h. The mixture was filtered and concentrated to give 501 mg (99 %) of the diphenylmethyl piperidine as a colorless oil.

- 4) TFA (1.4 ml) was added to a solution of the product of step 3 (500 mg, 1.4 mmol) in CH₂Cl₂ (15 ml). The solution was stirred at 25 °C for 23 h. The solution was concentrated and the residue partitioned between CH₂Cl₂ and 1 N NaOH. The aqueous layer was extracted with CH₂Cl₂, the combined organic layers were dried over Na₂SO₄, filtered and concentrated to obtain 349 mg (99 %) of the free amine as a yellow oil.

- 62 -

m.p. (HCl) = decomp. above 220-230 °C. HRMS calc'd for C₁₈H₂₂N (M⁺): 252.1752, Found: 252.1751.

- 5) A solution of the product of step 4 (349 mg, 1.4 mmol), N-Boc-4-piperidone 280 mg, 1.4 mmol), and Ti(OiPr)₄ (0.42 ml, 1.4 mmol) were taken up in CH₂Cl₂ (15 ml) under N₂. After stirring at 25 °C for 17 h, Et₃AlCN (2.8 mmol, 2.8 ml of 1.0 M in toluene) was added and the solution was stirred an additional 18 h at 25 °C. The solution was quenched with sat. NaHCO₃, diluted with EtOAc and filtered through Celite. The aqueous layer was extracted with EtOAc and the combined EtOAc layers were dried over Na₂SO₄. Filtration and concentration gave a yellow oil. Purification via preparative layer chromatography (3/1 hexanes/EtOAc, SiO₂) gave 430 mg (67 %) of the desired product as an oil.


- 6) A solution of the product of step 5 (430 mg, 0.94 mmol) in THF (20 ml) was cooled to 0 °C under N₂. CH₃MgBr (1.6 ml of 3.0 M in Et₂O, 4.7 mmol) was added at 0 °C and the solution stirred at 25 °C for 19 h. The reaction mixture was quenched with sat. NH₄Cl, diluted with CH₂Cl₂ and 1 N NaOH (check aqueous layer with pH paper, pH = 8-10). The layers were separated and the aqueous layer extracted with CH₂Cl₂. The combined organic layers were dried over Na₂SO₄, filtered and concentrated to obtain a yellow oil. Purification via flash chromatography (3/1 hexanes/EtOAc, SiO₂) gave 275 mg (65 %) of the product as a yellow oil.

- 7) TFA (0.60 ml) was added to a solution of the product of step 6 (275 mg, 0.61 mmol) in CH₂Cl₂ (15 ml) and the solution was stirred at 25 °C for 18 h. The solution was concentrated and the residue was partitioned between CH₂Cl₂ and 1 N NaOH. The aqueous layer was extracted with CH₂Cl₂, the combined organic layers were dried over Na₂SO₄, filtered and concentrated to obtain 209 mg (99 %) of the amine as a yellow oil. HRMS calc'd for C₂₄H₃₃N₂ (M⁺): 349.2644, Found: 349.2638.

- 8) A solution of the product of step 7 (50 mg, 0.14 mmol), 2,6-dimethylbenzoic acid (63 mg, 0.42 mmol), EDCI (54 mg, 0.28 mmol), HOBt (38 mg, 0.28 mmol), and IP₂NEt (0.10 ml) were taken up in CH₂Cl₂ (3 ml). The solution was stirred at 25 °C for 18 h, then diluted with CH₂Cl₂ and washed with 1 N NaOH. The aqueous layer was extracted with CH₂Cl₂, the combined organic layers were dried over Na₂SO₄, and filtered and concentrated to give a yellow oil. Purification via preparative thin-layer chromatography (3/1 hexanes/EtOAc, SiO₂) gave 47 mg (70 %) of the title

compound as a colorless oil, m.p. (HCl salt) = 195-201 °C. HRMS calc'd for C₃₃H₄₁N₂O (MH⁺): 481.3219, Found: 481.3225.



Using similar procedures, compounds of the following structure were prepared



wherein R⁶ and R² are as defined in the table:

Ex.	R ⁶	R ²	HRMS (MH ⁺) found	M.p., °C (HCl salt)
11A	H	H ₃ C-C ₆ H ₃ (CH ₃) ₂ -N	482.3156	201-207
11B	F ₃ CO-	H ₃ C-C ₆ H ₃ (CH ₃) ₂ -N	565.3069	204-209
11C	H	H ₃ C-C ₆ H ₃ (CH ₃) ₂ -N	482.3168	187-192
11D	F ₃ CO-	H ₃ C-C ₆ H ₃ (CH ₃) ₂ -N	567.2957	175-181
11E	F ₃ CO-	H ₃ C-C ₆ H ₃ (CH ₃) ₂ -N	582.2966	92-98
11F	F ₃ CO-	H ₃ C-C ₆ H ₃ (CH ₃) ₂ -N	566.3020	175-181

Example 12



- 1) N-Boc-4-piperidone (10 g, 50 mmol) and diethyl benzylphosphonate (12.6g, 55 mmol) were taken up in dry THF (50 ml) under N₂. NaH (2.4 g, 60 mmol, 60 wt % in oil dispersion) was added to the solution at 25 °C. The resulting mixture was heated at reflux for 3.5 h. The solution was partitioned between EtOAc and saturated NH₄Cl, the aqueous layer was extracted with EtOAc and the combined EtOAc layers were washed with brine and dried over MgSO₄. Filtration and concentration afforded a yellow oil. Purification via flash chromatography (10/1 hexanes/Et₂O, SiO₂) gave 9.85 g (72 %) of the desired compound as a solid, m.p. = 63-65 °C.
- 2) Bromine (1 ml, 20 mmol, dissolved in 10 ml CH₂Cl₂) was added dropwise to a CH₂Cl₂ (100 ml) solution of the product of step 1 (5.0 g, 18 mmol) at 0 °C. The solution was stirred at 0 °C for 15 min, then concentrated under reduced pressure. The crude product was taken up in tert-butanol/THF (4/1, 100 ml), and KOtBu (4.1 g, 36 mmol) was added to the solution in portions. The yellow mixture was stirred at 25 °C for 5h, then concentrated under reduced pressure. The residue was partitioned between EtOAc and saturated NH₄Cl, the aqueous layer was extracted with EtOAc, and the combined EtOAc layers were washed with brine and dried over MgSO₄. Filtration and concentration gave a yellow solid. Purification via flash chromatography (7/1 hexanes/Et₂O, SiO₂) gave 5.2 g (81 %) of the desired product as a yellow solid, m.p. = 80-83 °C.

- 65 -

- 3) TFA (5.9 ml) was added to a solution of the product of step 2 (2.1 g, 5.9 mmol) in CH_2Cl_2 (25 ml). The solution was stirred at 25 °C for 5 h, concentrated and the residue was partitioned between CH_2Cl_2 and 1 N NaOH. The aqueous layer was extracted with CH_2Cl_2 and the combined organic layers were dried over Na_2SO_4 , filtered and concentrated to obtain 1.46 g (98 %) of the amine as an orange oil, m.p. (HCl salt) = decomp. above 185-195 °C. HRMS calcd for $\text{C}_{12}\text{H}_{15}\text{BrN}$ (MH^+): 254.0367. Found: 254.0374.
- 4) A solution of the product of step 3 (1.4 g, 5.6 mmol), N-Boc-4-piperidone (1.1 g, 5.6 mmol), and $\text{Ti}(\text{O}i\text{Pr})_4$ (1.7 ml, 5.6 mmol) were taken up in CH_2Cl_2 (30 ml) under N_2 . After stirring at 25 °C for 18 h, Et_3AlCN (6.7 mmol, 6.7 ml, 1.0 M in toluene) was added to the solution and the solution was stirred an additional 18 h at 25 °C. The solution was quenched with sat. NaHCO_3 , diluted with EtOAc and filtered through Celite.
- 5) The aqueous layer was extracted with EtOAc and the combined EtOAc layers were dried over Na_2SO_4 . Filtration and concentration gave a yellow oil. Purification via flash chromatography (3/1 hexanes/EtOAc, SiO_2) gave 2.0 g (78 %) of the desired product as an off-white solid.
- 5) A solution of the product of step 4 (2.0 g, 4.3 mmol) in THF (30 ml) was cooled to 0 °C under N_2 . CH_3MgBr (7.2 ml of 3.0 M in Et_2O , 21 mmol) was added to the solution at 0 °C. The solution was warmed to 25 °C and stirred at that temperature for 16 h. The reaction mixture was quenched with sat. NH_4Cl and diluted with CH_2Cl_2 and 1 N NaOH (check aqueous layer with pH paper, pH = 8-10). The layers were separated, the aqueous layer was extracted with CH_2Cl_2 and the combined organic layers were dried over Na_2SO_4 . Filtration and concentration gave a yellow oil. Purification via flash chromatography (3/1 hexanes/EtOAc, SiO_2) gave 1.56 g (82 %) of the desired product as a yellow oil.
- 6) A solution of the product of step 5 (300 mg, 0.67 mmol), $4\text{-CF}_3\text{C}_6\text{H}_4\text{B}(\text{OH})_2$ (380 mg, 2 mmol), $\text{PdCl}_2(\text{PPh}_3)_2$ (50 mg, 0.067 mmol), and Na_2CO_3 (210 mg, 2 mmol) were taken up in THF/ H_2O (4/1, 15 ml) and stirred at 65 °C under N_2 for 18 h. The solution was partitioned between EtOAc and H_2O and the aqueous layer was extracted with EtOAc. The combined organic layers were washed with brine and dried over Na_2SO_4 . Filtration and concentration gave a dark brown oil. Purification via flash chromatography (4/1 hexanes/EtOAc, SiO_2) gave 229 mg (67 %) of the desired product as a colorless oil.

- 66 -

- 7) A solution of the product of step 6 (229 mg, 0.45 mmol) and $\text{Pd}(\text{OH})_2$ on carbon (200 mg, 20 wt % Pd (dry basis), 50 wt % H_2O) were taken up in CH_3OH (35 ml) and shaken in a Parr apparatus under H_2 (50 psi) for 20 h. The mixture was filtered and concentrated to obtain 232 mg (100 %) of the (±)-product as a colorless foam. HRMS calcd for $\text{C}_{30}\text{H}_{40}\text{O}_2\text{N}_3$ (MH^+): 517.3042. Found: 517.3050.
- 8) TFA (0.45 ml) was added to a solution of the product of step 7 (235 mg, 0.45 mmol) in CH_2Cl_2 (15 ml). The solution was stirred at 25 °C for 24 h, then concentrated and the residue was partitioned between CH_2Cl_2 and 1 N NaOH. The aqueous layer was extracted with CH_2Cl_2 , the combined organic layers were dried over Na_2SO_4 , filtered and concentrated to obtain 146 mg (78 %) of the (±)-amine as a yellow oil.
- 9) A solution of the product of step 8 (102 mg, 0.25 mmol), 4,6-dimethylpyrimidine-5-carboxylic acid (110 mg, 0.75 mmol), EDCI (96 mg, 0.50 mmol), HOBT (70 mg, 0.50 mmol), and Pr_2NEt (0.17 ml) was taken up in CH_2Cl_2 (3 ml). The solution was stirred at 25 °C for 18 h, then diluted with CH_2Cl_2 and washed with 1 N NaOH. The aqueous layer was extracted with CH_2Cl_2 , the combined organic layers were dried over Na_2SO_4 , filtered and concentrated to obtain a yellow oil. Purification via preparative thin-layer chromatography (1/1 acetone/hexanes SiO_2) gave 121 mg (88 %) of the title compound as a colorless oil, m.p. (HCl salt) = 186-191 °C. HRMS calcd for $\text{C}_{32}\text{H}_{38}\text{N}_4\text{O}_3$ (MH^+): 551.2998. Found: 551.3012.
- The 4,6-dimethylpyrimidine-5-carboxylic acid used in step 9 was made by the following process:
-
- Step 1: Ethyl diacetacetate (93.4 g), Cs_2CO_3 (155 g) and CH_3CN (550 ml) were mixed together, using an overhead mechanical stirrer. CH_3CN (50 ml) was added and the resulting mixture was cooled to 0 °C. Methyl trifluoromethane sulfonate (88.6 g) was added dropwise and after addition, the cooling bath was removed. The mixture was stirred for 1 h at RT, filtered, and the salts were washed with Et_2O (2 X 50 ml). The organic extracts were combined and Et_2O (300 ml) was added. The resulting mixture was filtered, the filter cake was washed with Et_2O (2 X 100 ml), the Et_2O extracts were combined and evaporated to half volume. The solution was cooled in an ice bath and washed once with cooled (0 °C) 2 N NaOH

- 67 -


(pH = 11). The Et₂O layer was dried over MgSO₄, filtered and evaporated to give the desired product as a yellow liquid (64.7 g) in 65% yield, which was used directly in the next step.

- 5 **Step 2:** The product of step 1 (64.2 g), sodium ethoxide in ethanol (commercial solution; 21 wt%, 113 g) and formamidine acetate (36.2 g) were mixed together at RT. After refluxing for 4 h, the mixture was cooled to RT, the resulting precipitate was filtered off and the ethanol was removed under vacuum. The resulting liquid was partitioned between water and CH₂Cl₂ and the aqueous layer was extracted with CH₂Cl₂ (3 x 150 ml). The CH₂Cl₂ extracts were dried over MgSO₄, filtered and evaporated to give a dark crude liquid (50.7 g) which was purified by silica gel chromatography (980 g; 4:1 hexanes:EtOAc as eluant). After evaporation of the appropriate fractions, the desired product (28.5 g) was isolated in 46% yield and used directly in the next step.

- 15 **Step 3:** The product of step 2 (28.1 g), NaOH (6.72 g), water (65 ml) and EtOH (130 ml) were mixed together at RT and heated at reflux for 1h. The resulting solution was cooled to RT and the volatile materials were removed in vacuo until a thick paste resulted. Water (20 ml) was added, the mixture was cooled to 0°C and conc. HCl (14.3 ml) was added dropwise with stirring. The resulting white precipitate was collected by filtration, washed with ice water (2 X 10 ml) and air dried with suction for 30 min. The resulting white solid was treated with toluene (2 x 20 ml), the solvent was removed in vacuo at 50°C and then dried under vacuum (1 mm Hg) for 18 h. The desired product (14.9 g) was isolated as a white solid in 83% yield, mp: 176-178°C. Elemental analysis of C₂₇H₂₈N₂O₂: calcd C 55.26%, H 5.30%, N 18.41%; found: C 55.13%, H 5.44%, N 18.18%.

- 20 A second crop of product was isolated by evaporation of the aqueous filtrate (from above) to dryness and addition of water (20 ml). The resulting mixture was stirred at RT for 5 min, cooled in an ice bath and the precipitate formed was collected by filtration. The resulting solid was washed with ice water (2 X 5 ml) and dried as described above to give the product (4.68 g) as a cream colored solid to give a combined yield of 83%.

- 68 -

Example 13

- 5 **Step 1:** To a suspension of methyltriphenylphosphonium bromide (1.89 g; 4.80 mmol) in anhydrous THF (15 ml) at -40 °C is added n-BuLi 2.5 N in hexanes (2.12 ml; 5.3 mmol) via syringe. The reaction is allowed to warm to 0 °C, stirred 30 min at this temperature, and a solution of the product of Example 6, step B-2 (2.24 g; 4.8 mmol) is added. The solution is then allowed to warm to RT overnight, poured into CH₂Cl₂ and washed with saturated NaHCO₃ then brine. The residue obtained after concentration of the organic layer is purified by flash chromatography over silica gel (eluting with CH₂Cl₂/EtOAc, 9:1) to afford 0.56 g (25%) of an oil.



- 10 **Step 2:** A solution of the product of step 1 (0.56 g; 1.2 mmol) and 9-BBN, 0.5 N in THF (3 ml; 1.5 mmol) is refluxed 2 h under inert atmosphere. Part of this solution (1.5 ml; 0.59 mmol of theoretical intermediate) is added to a mixture of 1-chloro-3-iodobenzene (88 µl; 0.71 mmol), PdCl₂(dppf).CH₂Cl₂ (19.8 mg), triphenylarsine (24.1 mg) and Cs₂CO₃ (250 mg) in DMF (0.40 ml) and water (80 µl). The reaction is stirred 2 h at 60 °C and overnight at RT, poured into 5% aqueous NaHCO₃ and extracted with CH₂Cl₂. Combined organic layers are dried over Na₂SO₄, concentrated, and purified by chromatography over silica gel (eluting with EtOAc/hexanes, 8:2) to provide 100 mg (29%) of an oil.

- 15 **Step 3:** The Boc-protecting group of the product of step 2 (100 mg; 0.17 mmol) was removed as in Example 2 to obtain the desired amine (70 mg; 86%). This amine (45 mg; 0.09 mmol) was coupled with 4,6-dimethylpyrimidine-5-carboxylic acid following the conditions described in Example


- 69 -

- 2 to obtain the title compound as a colorless oil (32 mg). ¹H-NMR (300 MHz, CDCl₃) δ 8.93 (d, J = 3.8 Hz, 1H), 6.90-7.10 (m, 5H), 6.88 (br s, 1H), 6.71 (d, J = 7 Hz, 1H), 4.20 (m, 1H), 3.25-3.55 (m, 2H), 3.19 (m, 2H), 2.50-3.10 (m, 5H), 2.47 and 2.48 (s, 3H), 2.42 and 2.43 (s, 3H), 1.70-2.20 (m, 5H), 1.20-1.65 (m, 5H), 0.92 (s, 3H); HRMS (MH⁺) 615.2722.

Using a similar procedure, the following compound was also prepared:

13A HRMS (MH⁺) 599.3015**Example 14**

10



As in example 11, steps 7-8

As in example 11, steps 7-8

Enantiomer 2

To prepare a compound wherein R² is 2,6-dimethylphenyl:

- 1) A solution of the product of step 5 in example 12 (300 mg, 0.67 mmol), 4-CF₃OC₆H₄B(OH)₂ (410 mg, 2 mmol), PdCl₂(PPh₃)₂ (50 mg, 0.067


- 70 -

- mmol), and Na₂CO₃ (210 mg, 2 mmol) were taken up in THF/H₂O (4/1, 15 ml) and stirred at 65 °C under N₂ for 19 h. The solution was partitioned between EtOAc and H₂O, and the aqueous layer was extracted with EtOAc. The combined organic layers were washed with brine and dried over Na₂SO₄. Filtration and concentration gave a dark brown oil. Purification via flash chromatography (4/1 hexanes/Et₂O, SiO₂) gave 356 mg (100 %) of the desired product as a yellow oil.










- 2) A solution of the product in step 1 (340 mg, 0.64 mmol) and Pd(OH)₂ on carbon (300 mg, 20 wt % Pd (dry basis), 50 wt % H₂O) were taken up in CH₃OH (35 ml) and shaken in a Parr apparatus under H₂ (50 psi) for 18 h. The mixture was filtered and concentrated to obtain 341 mg (100 %) of the product, (±)-1, as a colorless foam.

- 3) The amine (±)-1 was resolved via chiral HPLC separation. The conditions are as follows: CHIRALCEL® OD™ (5 cm x 30 cm); Hexane/isopropyl alcohol/diethylamine 75/25/0.05) at 25 °C; 254 nm detection. The retention times for peak 1, (+)-enantiomer, and peak 2, (-)-enantiomer were 3.8 and 4.9 minutes, respectively [CHIRALCEL® OD™ (hexane/ethanol/diethylamine 90/10/0.1) 25 °C at 254 nm]. Peak 1 and peak 2 are the first and second eluting peaks from the column, respectively. The enantiomers (I and II) were deprotected (CH₂Cl₂/TFA), and the free amine was coupled to the 2,6-dimethylbenzoic acid using the conditions described in example 11, steps 7 and 8. The hydrochloride salts were obtained by taking the free base up in EtOAc and titrating with 1 M HCl in Et₂O.



- Data for the above compounds, 14A and 14B, and for additional compounds made in a similar manner, are given in the following table. In each case, the enantiomer designator I is derived from (+)-1 and the enantiomer designated II is derived from (-)-1.



Ex.	Ar	Enantiomer	m.p. (HCl)	HRMS	
				calc	found
14A		I	185-190	565.3042	565.3050

14B		II	175-180	565.3042	566.3050
14C		I	168-174	567.2947	567.2951
14D		II	170-175	567.2947	567.2957
14E		I	195-201	582.2944	582.2944
14F		II	180-185	582.2944	582.2958
14G		II	214-218	581.2991	581.2984
14H		II	145-151	658.3257	658.3251
14I		II	193-198	615.3010	615.3016
14J		II	195-200	651.3522	651.3526

Example 15



1) The dibromo-olefin (3.55 g, 10 mmol) and TFA (10 ml) were taken up in CH_2Cl_2 and stirred at 25 °C for 20 h. The solution was concentrated. The residue was partitioned between CH_2Cl_2 and 1 N NaOH. The aqueous layer was extracted with CH_2Cl_2 . The combined organic layers were dried (Na_2SO_4). Filtration and concentration gave the 2.4 g (94 %) of the free piperidine as a colorless oil. The free piperidine (2.41 g, 9.45 mmol) was treated sequentially with (a) N-Boc-4-piperidone/Tf(OiPr)₃ and (b) Et_3AlCl , to give the cyano-amine as described in Step 5 of Example 11.


10 2) The product of Step 1 and MeMgBr (16 ml, 3.0 M in Et_2O) were taken up in THF (30 ml) and stirred at 25 °C for 19 h. The solution was quenched with 1 N NaOH and EtOAc. The mixture was filtered (Celite). The aqueous layer was extracted with EtOAc, the combined EtOAc layers were washed with brine and dried (Na_2SO_4). Filtration and concentration gave a yellow oil. Purification via flash chromatography (6/1 hexanes/EtOAc, SiO_2) gave 2.54 g (69 % from the free piperidine) of the vinyl bromide as a solid. m.p. (free base) 85–90 °C. HRMS (MH^+) calcd. for $\text{C}_{10}\text{H}_{12}\text{O}_2\text{N}_2\text{Br}$, 387.1647; Found, 387.1638.

3) The product of Step 2 (200 mg, 0.52 mmol), 4- $\text{CF}_3\text{C}_6\text{H}_4\text{B}(\text{OH})_2$ (344 mg, 1.8 mmol), $\text{PdCl}_2(\text{PPh}_3)_2$ (36 mg, 0.52 mmol), and Na_2CO_3 (165 mg, 1.55 mmol) were taken up in $\text{THF}/\text{H}_2\text{O}$ (4/1, 10 ml) and heated at 75 °C (oil bath) for 21 hours. The solution was partitioned between EtOAc and H_2O . The aqueous layer was extracted with EtOAc, the combined EtOAc layers were washed with brine and dried (Na_2SO_4). Filtration and concentration gave a yellow oil. Purification via flash chromatography (3/1 to 1/1 hexanes/EtOAc, SiO_2) gave 210 mg (89 %) of the phenyl substituted olefin as an oil. HRMS (M^+) calcd. for $\text{C}_{26}\text{H}_{19}\text{O}_2\text{NF}_3$, 453.2729; Found, 453.2728.

4) The product of Step 3 was hydrogenated as described in Step 3 of Example 11. The reduced product was deprotected and coupled to 2,6-dimethyl benzoic acid as described in Example 11, steps 7-8 to give the title compound as a yellow oil (37 mg, 55%), m.p. (HCl salt) 130-140 °C. HRMS (MH⁺) calcd. for C₂₈H₃₈ON₂F₅, 487.2936; Found, 487.2928.

Using a similar procedure, the following compound was prepared:

- 73 -



m.p. (HCl salt) 135-145°C. HRMS (MH⁺) calcd. for C₂₂H₂₀O₄N₂F₃, 503.2885; Found, 503.2896.

5 The following assays can be used to determine the CCR5 inhibitory and antagonistic activity of the compounds of the invention.

CCR5 Membrane Binding Assay:

A high throughput screen utilizing a CCR5 membrane binding assay identifies inhibitors of RANTES binding. This assay utilizes membranes prepared from NIH 3T3 cells expressing the human CCR5 chemokine receptor which have the ability to bind to RANTES, a natural ligand for the receptor. Using a 96-well plate format, membrane preparations are incubated with 125I-RANTES in the presence or absence of compound for one hour. Compounds are serially diluted over a wide range of 0.001 µg/ml to 1 µg/ml and tested in triplicates. Reaction cocktails are harvested through glass fiber filters, and washed thoroughly. Total counts for replicates are averaged and data reported as the concentration required to inhibit 50 percent of total 125I-RANTES binding. Compounds with potent activity in the membrane binding assay are further characterized in secondary cell-based HIV-1 entry and replication assays.

HIV-1 Entry Assay:

Replication defective HIV-1 reporter virions are generated by cotransfection of a plasmid encoding the NL4-3 strain of HIV-1 (which has been modified by mutation of the envelope gene and introduction of a luciferase reporter plasmid) along with a plasmid encoding one of several HIV-1 envelope genes as described by Connor et al., Virology, 206 (1995), p. 935-944. Following transfection of the two plasmids by calcium phosphate precipitation, the viral supernatants are harvested on day 3 and a functional viral titer determined. These stocks are then used to infect U87 cells stably expressing CD4 and the chemokine receptor CCR5 which have been preincubated with or without test compound. Infections are carried out for 2 hours at 37 °C, the cells washed and media replaced with fresh media containing compound. The cells are incubated for 3 days, lysed and luciferase

- 74 -

activity determined. Results are reported as the concentration of compound required to inhibit 50% of the luciferase activity in the control cultures.

HIV-1 Replication Assay:

This assay uses primary peripheral blood mononuclear cells or the stable U87-CCR5 cell line to determine the effect of anti-CCR5 compounds to block infection of primary HIV-1 strains. The primary lymphocytes are purified from normal healthy donors and stimulated *in vitro* with PHA and IL-2 three days prior to infection. Using a 96-well plate format, cells are pretreated with drug for 1 hour at 37 °C and subsequently infected with an M-tropic HIV-1 isolates. Following infection, the cells are washed to remove residual inoculum and cultured in the presence of compound for 4 days. Culture supernatants are harvested and viral replication measured by determination of viral p24 antigen concentration.

Calcium Flux Assay:

Cells expressing the HIV coreceptor CCR5 are loaded with calcium sensitive dyes prior to addition of compound or the natural CCR5 ligand. Compounds with agonist properties will induce a calcium flux signal in the cell, while CCR5 antagonists are identified as compounds which do not induce signaling by themselves but are capable of blocking signaling by the natural ligand RANTES.

GTPγS Binding Assay (secondary membrane binding assay):

A GTPγS binding assay measures receptor activation by CCR5 ligands. This assay measures the binding of 35S labeled-GTP to receptor coupled G-proteins that occurs as a result of receptor activation by an appropriate ligand. In this assay, the CCR5 ligand, RANTES, is incubated with membranes from CCR5 expressing cells and binding to the receptor activation (or binding) is determined by assaying for bound 35S label. The assay quantitatively determines if compounds exhibit agonist characteristics by inducing activation of the receptor or alternatively antagonist properties by measuring inhibition of RANTES binding in a competitive or non-competitive fashion.

Chemotaxis Assay:

The chemotaxis assay is a functional assay which characterizes the agonist vs. antagonist properties of the test compounds. The assay measures the ability of a non-adherent murine cell line expressing human CCR5 (BaF-550) to migrate across a membrane in response to either test compounds or natural ligands (i.e., RANTES, MIP-1β). Cells migrate across the permeable membrane towards compounds with agonist activity. Compounds that are antagonists not only fail to induce chemotaxis, but are

- 75 -

also capable of inhibiting cell migration in response to known CCR5 ligands.

The role of CC chemokine receptors such as CCR-5 receptors in inflammatory conditions has been reported in such publications as

- 5 Immunology Letters, 57, (1997), 117-120 (arthritis); Clinical & Experimental Rheumatology, 17 (4) (1999), p. 419-425 (rheumatoid arthritis); Clinical & Experimental Immunology, 117 (2) (1999), p.237-243 (atopic dermatitis); International Journal of Immunopharmacology, 20 (11) (1998), p. 661-7 (psoriasis); Journal of Allergy & Clinical Immunology, 100 (6, Pt 2) (1997), p. S52-5 (asthma); and Journal of Immunology, 159 (6) (1997), p. 2962-72 (allergies).

- 10 In the assay to determine inhibition of RANTES binding, compounds of the invention range in activity from a K_i of 0.1 to 2000 nM, with preferred compounds having a range of activity from 0.1 to 1000 nM, more preferably 0.1 to 500 nM, and most preferably 0.1 to 100 nM. The results for

15 preferred and representative compounds of formulas I and II in the test to determine inhibition of RANTES binding are given in the table below. In the table, "Ex. No." stands for "Example Number" and "nM" stands for "nanomolar."

Ex. No.	K_i (nM) Inhibition of RANTES binding
1B	14
1J	1
2	9.6
2G	1.8
2S	17.9
2LJ	0.58
4B	0.5
4C	0.5
5L	7.9
5N	1.7
5O	0.4
5Z	0.3
5AB	0.1
6V	0.8
7U	62.5
9D	588

- 76 -

For preparing pharmaceutical compositions from the CCR5

- 5 antagonist compounds described by this invention, inert, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, dispersible granules, capsules, cachets and suppositories. The powders and tablets may be comprised of from about 5 to about 95 percent active ingredient. Suitable solid carriers are known in the art, e.g., magnesium carbonate, magnesium stearate, talc, sugar or lactose. Tablets, powders, cachets and capsules can be used as solid
- 10 dosage forms suitable for oral administration. Examples of pharmaceutically acceptable carriers and methods of manufacture for various compositions may be found in A. Gennaro (ed.), Remington's Pharmaceutical Sciences, 18th Edition, (1990), Mack Publishing Co., Easton, Pennsylvania.

- 15 Liquid form preparations include solutions, suspensions and emulsions. As an example may be mentioned water or water-propylene glycol solutions for parenteral injection or addition of sweeteners and opacifiers for oral solutions, suspensions and emulsions. Liquid form preparations may also include solutions for intranasal administration.
- 20 Aerosol preparations suitable for inhalation may include solutions and solids in powder form, which may be in combination with a pharmaceutically acceptable carrier, such as an inert compressed gas, e.g., nitrogen.

- 25 Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms include solutions, suspensions and emulsions.

- 30 The compounds of the invention may also be deliverable transdermally. The transdermal compositions can take the form of creams, lotions, aerosols and/or emulsions and can be included in a transdermal patch of the matrix or reservoir type as are conventional in the art for this purpose.

Preferably the compound is administered orally.

- 35 Preferably, the pharmaceutical preparation is in a unit dosage form. In such form, the preparation is subdivided into suitably sized unit doses containing appropriate quantities of the active component, e.g., an effective amount to achieve the desired purpose.

- 77 -

The quantity of active compound in a unit dose of preparation may be varied or adjusted from about 10 mg to about 500 mg, preferably from about 25 mg to about 300 mg, more preferably from about 50 mg to about 250 mg, and most preferably from about 55 mg to about 200 mg, according to the particular application.

The actual dosage of CCR5 compound employed may be varied depending upon the requirements of the patient and the severity of the condition being treated. Determination of the proper dosage regimen for a particular situation is within the skill of the art. For convenience, the total daily dosage may be divided and administered in portions during the day as required.

The amount and frequency of administration of the CCR5 compounds of the invention and/or the pharmaceutically acceptable salts thereof will be regulated according to the judgment of the attending clinician considering such factors as age, condition and size of the patient as well as severity of the symptoms being treated. A typical recommended daily dosage regimen for oral administration can range from about 100 mg/day to about 300 mg/day, preferably 150 mg/day to 250 mg/day, more preferably about 200 mg/day, in two to four divided doses.

The doses and dosage regimens of the NRTIs, NNRTIs, PIs and other agents used in combination with the CCR5 antagonists will be determined by the attending clinician in view of the approved doses and dosage regimens in the package inserts or as set forth in the protocols, taking into consideration the age, sex and condition of the patient and the severity of the condition treated.


The goal of the HIV-1 therapy of the present invention is to reduce the HIV-1-RNA viral load below the detectable limit. The "detectable limit of HIV-1-RNA" in the context of the present invention means that there are fewer than about 200 to fewer than about 50 copies of HIV-1-RNA per ml of plasma of the patient as measured by quantitative, multi-cycle reverse transcriptase PCR methodology. HIV-1-RNA is preferably measured in the present invention by the methodology of Amplicor -1 Monitor 1.5 (available from Roche Diagnostics) or of Nuclisens HIV-1 QT -1.

While the present invention has been described in conjunction with the specific embodiments set forth above, many alternatives, modifications and variations thereof will be apparent to those of ordinary skill in the art. All such alternatives, modifications and variations are intended to fall within the spirit and scope of the present invention.

- 78 -

WHAT IS CLAIMED IS:

1. A compound represented by the structural formula II



or a pharmaceutically acceptable salt thereof, wherein
(1) X^8 is $-C(R^{13})_2$, $-C(R^{13})(R^{19})$, $-C(O)-$, $-O-$, $-NH-$, $-N((C_1-C_6)alkyl)$,

OR^3 , $-CH_2-(C_1-C_6)alkyl-R^3$, $-NOR^4$, $-O-(C_1-C_6)alkyl$, $-CH-(C_1-C_6)alkyl$,
 $-CR^{13}_2$, $-CR^{13}_-$, $-C-$, $-CR^{13}_-$, $-C-$

$-O-C(O)-(C_1-C_6)alkyl$, $-O-C(O)-O-(C_1-C_6)alkyl$, $-O-C(O)-NH-(C_1-C_6)alkyl$,
 $-CR^{13}_-$, $-CR^{13}_-$, $-CR^{13}_-$

$-O-C(O)-N((C_1-C_6)alkyl)_2$, $-NR^5-C(O)-(C_1-C_6)alkyl$,
 $-CR^{13}_-$, $-CR^{13}_-$

$-NR^5-C(O)-O-(C_1-C_6)alkyl$, $-NR^5-C(O)-NH-(C_1-C_6)alkyl$,
 $-CR^{13}_-$, $-CR^{13}_-$


$-NR^5-C(O)-N((C_1-C_6)alkyl)_2$, $-C(O)-(C_1-C_6)alkyl$,
 $-CR^{13}_-$, or $-N-$

R^9 is R^{6a} -phenyl, R^{6a} -pyridyl, R^{6a} -thiophenyl or R^6 -naphthyl;

R^1 is hydrogen, C_1-C_6 alkyl or C_2-C_6 alkenyl;

R^2 is R^7 , R^8 , R^9 -phenyl; R^7 , R^8 , R^9 -substituted 6-membered heteroaryl N-oxide;

heteroaryl; R^7 , R^8 , R^9 -substituted 6-membered heteroaryl N-oxide;
 R^{10} , R^{11} -substituted 5-membered heteroaryl; naphthyl; fluorenyl;



R^3 is R^{10} -phenyl, pyridyl, pyrimidyl, pyrazinyl or thiazolyl;

R^4 is hydrogen, C_1-C_6 alkyl, fluoro- C_1-C_6 alkyl, cyclopropylmethyl,


$-CH_2CH_2OH$, $-CH_2CH_2-O-(C_1-C_6)alkyl$, $-CH_2C(O)-O-(C_1-C_6)alkyl$,

$-CH_2C(O)NH_2$, $-CH_2C(O)-NH(C_1-C_6)alkyl$ or $-CH_2C(O)-N((C_1-C_6)alkyl)_2$;

R^5 and R^{11} are independently selected from the group consisting of hydrogen and (C_1-C_6) -alkyl;

- 79 -

R_{6a} is 1 to 3 substituents independently selected from the group consisting of hydrogen, halogen, -CF₃, CF₃O-, -CN, -CF₃SO₂, R²-phenyl,

-NHCOCF₃, 5-membered heteroaryl and , wherein X is -O-, -NH- or -N(CH₃)₂;

5 R₆ is independently selected from the group consisting of R_{6a} and CH₃SO₂-;

R₇ and R₈ are independently selected from the group consisting of (C₁-C₆)alkyl, halogen, -NR^aR^b, -OH, -CF₃, -OCH₃, -O-acyl, and -OCF₃;

R₉ is Rⁱ, hydrogen, phenyl, -NO₂, -CN, -CH₂F, -CHF₂, -CHO,

10 -CH=NO^a, pyridyl, pyridyl N-oxide, pyrimidinyl, pyrazinyl,

-N(R²⁰)CONR²¹R²², -NHCONH(chloro-(C₁-C₆)alkyl), -NHCONH((C₃-C₁₀)-

cycloalkyl)(C₁-C₆)alkyl), -NHCO(C₁-C₆)alkyl, -NHCOCF₃, -NHSO₂N((C₁-

C₆)alkyl)₂, -NHSO₂(C₁-C₆)alkyl, -N(SO₂CF₃)₂, -NHCO₂(C₁-C₆)alkyl, C₃-C₁₀

cycloalkyl, -SR²³, -SOR²³, -SO₂R²³, -SO₂NH(C₁-C₆)alkyl), -OSO₂(C₁-

15 C₆)alkyl, -OSO₂CF₃, hydroxy(C₁-C₆)alkyl, -CON R²⁰R²¹, -CON(CH₂CH₂O-CH₃)₂,

-OCONH(C₁-C₆)alkyl, -CO₂R²⁰, -Si(CH₃)₃ or -B(OC(CH₃)₂)₂;

R¹⁰ is (C₁-C₆)alkyl, -NH₂ or R¹²-phenyl;

R¹² is 1 to 3 substituents independently selected from the group

20 consisting of hydrogen, (C₁-C₆) alkyl, -CF₃, -CO₂R₂₀, -CN, (C₁-C₆)alkoxy and halogen;

R¹³, R¹⁴, R¹⁵ and R¹⁶ are independently selected from the group

consisting of hydrogen and (C₁-C₆)alkyl;

R¹⁷ and R¹⁸ are independently selected from the group consisting of

25 hydrogen and C₁-C₆ alkyl, or R¹⁷ and R¹⁸ together are a C₂-C₅ alkylene group and with the carbon to which they are attached form a spiro ring of 3 to 6 carbon atoms;

R¹⁹ is R⁵-phenyl, R⁵-heteroaryl, R⁵-naphthyl, C₃-C₁₀ cycloalkyl, (C₃-

30 C₁₀)cycloalkyl(C₁-C₆)alkyl or (C₁-C₆)alkoxy(C₁-C₆)alkyl;

R²⁰, R²¹ and R²² are independently selected from the group

consisting of H and C₁-C₆ alkyl; and

R²³ is C₁-C₆ alkyl or phenyl; or

(2):

X^a is -C(R¹³)(R¹⁹)-, -C(O)-, -O-, -NH-, -N((C₁-C₆)alkyl)-,

35

- 80 -

OR³ CH₂(C₁-C₅)alkyl-R³ NOR^{4a} O-C(O)-(C₁-C₆)alkyl

-CR¹³-, -CR¹³-, -C-, -CR¹³-

O-C(O)-(C₁-C₆)alkyl O-C(O)-NH-(C₁-C₆)alkyl

-CR¹³-, -CR¹³-

5 O-C(O)-N((C₁-C₆)alkyl)₂ NR⁵-C(O)-(C₁-C₆)alkyl

-CR¹³-, -CR¹³-

NR⁵-C(O)-O-(C₁-C₆)alkyl NR⁵-C(O)-NH-(C₁-C₆)alkyl

-CR¹³-, -CR¹³-

NR⁵-C(O)-N((C₁-C₆)alkyl)₂ Q(O)-(C₁-C₆)alkyl

-CR¹³-, or -N-

10 R_a is R^{6b}-phenyl, R^{6b}-pyridyl or R^{6b}-thiophenyl;

R^{4a} is fluoro-C₁-C₆ alkyl, cyclopropylmethyl, -CH₂CH₂OH,

-CH₂CH₂O-(C₁-C₆)alkyl, -CH₂C(O)-O-(C₁-C₆)alkyl, -CH₂C(O)NH₂, -

CH₂C(O)-NH-(C₁-C₆)alkyl or -CH₂C(O)-N((C₁-C₆)alkyl)₂;

R^{6b} is CH₃SO₂-; and

15 R¹, R², R³, R⁵, R¹⁴, R¹⁵, R¹⁶ and R¹⁹ are as defined in (1).

2. The compound of claim 1 wherein R_a is .

3. The compound of claim 1, formula II(1), wherein X^a is -CHOR³,

20 -C(R¹³)(R¹⁹)- or -C(=NOR^{4a})-.

4. The compound of claim 3 wherein R³ is pyridyl, R⁴ is (C₁-C₆)alkyl, or R³ is hydrogen and R¹⁹ is R⁵-phenyl.

25 5. The compound of claim 1, formula II(2), wherein X^a is -CHOR³, -C(R¹³)(R¹⁹)- or -C(=NOR^{4a})-.


6. The compound of claim 5 wherein R³ is pyridyl, R^{4a} is cyclopropylmethyl or trifluoroethyl, or R³ is hydrogen and R¹⁹ is R⁵-phenyl.

30

7. The compound of claim 1 wherein R² is R⁷, R⁸, R⁹-phenyl, R⁷, R⁸, R⁹-pyridyl or an N-oxide thereof, or R⁷, R⁸, R⁹-pyrimidinyl.


- 81 -

8. The compound of claim 7 wherein R² is selected from the group consisting of



- 5 wherein R⁷ and R⁸ are selected from the group consisting of (C₁-C₆)alkyl, halogen, and -NH₂, and R⁹ is hydrogen.

9. A compound of claim 1 selected from the group consisting of those represented by the formula



10

wherein R⁶, X and R² are as defined in the following table:

R ⁶	X	R ²
Br	OCH_2CH_3 -CH-	
Br	$\text{OCH}_2\text{CH}_2\text{CH}_3$ -CH-	
Br	$\text{OCH}_2\text{CH}_2\text{CH}_3$ -CH-	
Br	$\text{OCH}_2\text{CH}_2\text{CH}_3$ -CH-	
Br	$\text{OCH}_2\text{CH}_2\text{CH}_3$ -CH-	
Br	$\text{OCH}_2\text{CH}_2\text{CH}_3$ -CH-	
Br	$\text{OCH}_2\text{CH}_2\text{CH}_3$ -CH-	
Br	$\text{OCH}_2\text{CH}_2\text{CH}_3$ -CH-	
Br	$\text{OCH}_2\text{CH}_2\text{CH}_3$ -CH-	
Br	$\text{OCH}_2\text{CH}_2\text{CH}_3$ -CH-	

- 82 -

Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		

- 83 -

F ₃ CO-		
Br		
Br		
F ₃ CO-		
Br		
Br		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		
H ₃ CSO ₂ -		














Cl		
F		
Br		
Br		
Br		
Br		
F ₃ C-		
F ₃ C-		
F ₃ C-		
F ₃ C-		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		

- 87 -

Cl		
Cl		
F ₃ CO-		
Br		
Br		
Br		
Br		
Br		
Br		
Br		

- 88 -

Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		
Br		

Br	-CH ₂ -	
Br	-CH ₂ -	
Br	-CH ₂ -	
Br	-CH ₂ -	
Br	-CH ₂ -	
Br	-CH ₂ -	
Br	-CH-	
CH ₃ SO ₂ -	-CH-	
Br	-CH-	
Br	-CH-	
F	-CH-	
F	-CH-	
F	-CH-	

Br		
Cl		
F ₃ C-		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		
CH ₃ SO ₂ -		
F ₃ CO-		

- 99 -

- $$\begin{array}{c} \text{O}-\text{C}(\text{O})-\text{N}((\text{C}_1-\text{C}_6\text{alkyl})_2) \\ | \\ -\text{CR}^{13}- \end{array}$$

$$\begin{array}{c} \text{NR}^5-\text{C}(\text{O})-(\text{C}_1-\text{C}_6\text{alkyl}) \\ | \\ -\text{CR}^{13}- \end{array}$$


$$\begin{array}{c} \text{NR}^5-\text{C}(\text{O})-\text{O}-(\text{C}_1-\text{C}_6\text{alkyl}) \\ | \\ -\text{CR}^{13}- \end{array}$$

$$\begin{array}{c} \text{NR}^5-\text{C}(\text{O})-\text{NH}-(\text{C}_1-\text{C}_6\text{alkyl}) \\ | \\ -\text{CR}^{13}- \end{array}$$



$$\begin{array}{c} \text{NR}^5-\text{C}(\text{O})-\text{N}((\text{C}_1-\text{C}_6\text{alkyl})_2) \\ | \\ -\text{CR}^{13}- \end{array}$$

$$\begin{array}{c} \text{C}(\text{O})-(\text{C}_1-\text{C}_6\text{alkyl}) \\ | \\ -\text{N}- \end{array}$$
- 5 R is R⁶-phenyl, R⁶-pyridyl, R⁶-thiophenyl or R⁶-naphthyl;
 R¹ is hydrogen, C₁-C₆ alkyl or C₂-C₆ alkenyl;
 R² is R⁷, R⁸, R⁹-phenyl; R⁷, R⁸, R⁹-substituted 6-membered heteroaryl; R⁷, R⁸, R⁹-substituted 6-membered heteroaryl N-oxide;
- 10 R¹⁰, R¹¹-substituted 5-membered heteroaryl; naphthyl; fluorenyl;
- $$\begin{array}{c} \text{R}^{17} \\ | \\ \text{C} \\ | \\ \text{R}^{18} \end{array}$$

$$\begin{array}{c} \text{R}^{17} \\ | \\ \text{C} \\ | \\ \text{R}^{18} \end{array}$$

$$\begin{array}{c} \text{R}^{17} \\ | \\ \text{C} \\ | \\ \text{R}^{18} \end{array}$$
- R³ is R⁶-phenyl, R⁴-heteroaryl or R⁵-naphthyl;
 R⁴ is hydrogen, C₁-C₆ alkyl, fluoro-C₁-C₆ alkyl, cyclopropylmethyl, -CH₂CH₂OH, -CH₂CH₂-O-(C₁-C₆alkyl), -CH₂C(O)-O-(C₁-C₆alkyl), -CH₂C(O)NH₂, -CH₂C(O)-NH(C₁-C₆alkyl) or -CH₂C(O)N(C₁-C₆alkyl)₂;
- 15 R⁵ and R¹¹ are independently selected from the group consisting of hydrogen and (C₁-C₆)-alkyl;
- R⁶ is 1 to 3 substituents independently selected from the group consisting of hydrogen, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, -CF₃, CF₃O-, CH₃C(O)-, -CN, CH₃SO₂-, CF₃SO₂-, R¹⁴-phenyl, R¹⁴-benzyl,
- 20 CH₃C(=NOCH₃), CH₃C(=NOCH₂CH₃), NHCONH(C₁-C₆ alkyl), -NHCO(C₁-C₆ alkyl), -NHSO₂(C₁-C₆ alkyl),
- 25 5-membered heteroaryl and , wherein X is -O-, -NH- or -N(CH₃)₂; R⁷ and R⁸ are independently selected from the group consisting of (C₁-C₆alkyl), halogen, -NR²⁰R²¹, -OH, -CF₃, -OCH₃, -O-acyl and -OCF₃; R⁹ is R⁷, hydrogen, phenyl, -NO₂, -CN, -CH₂F, -CHF₂, -CHO, -CH=NOR²⁰, pyridyl, pyridyl N-oxide, pyrimidinyl, pyrazinyl, -N(R²⁰)CONR²¹R²², -NHCONH(chloro-(C₁-C₆alkyl)), -NHCONH((C₂-C₆)-cycloalkyl(C₁-C₆alkyl)), -NHCO(C₁-C₆alkyl), -NHCOCF₃, -NHSO₂N((C₁-C₆alkyl)₂), -NHSO₂(C₁-C₆alkyl), -N(SO₂CF₃)₂, -NHSO₂(C₁-C₆alkyl), C₃-C₆

- 100 -

- cycloalkyl, -SR²³, -SOR²³, -SO₂R²³, -SO₂NH(C₁-C₆ alkyl), -OSO₂(C₁-C₆alkyl), -OSO₂CF₃, hydroxy(C₁-C₆alkyl), -CON R²⁴R²⁵, -CON(CH₂CH₂-O-CH₃)₂, -OCONH(C₁-C₆alkyl), -CO₂R²⁶, -Si(CH₃)₃ or -B(OC(CH₃)₂)₂;
- 5 R¹⁰ is (C₁-C₆alkyl), -NH₂ or R¹²-phenyl;
- R¹² is 1 to 3 substituents independently selected from the group consisting of hydrogen, (C₁-C₆ alkyl), -CF₃, -CO₂R²⁶, -CN, (C₁-C₆alkoxy and halogen;
- 10 R¹³, R¹⁴, R¹⁵ and R¹⁶ are independently selected from the group consisting of hydrogen and (C₁-C₆alkyl);
- R¹⁷ and R¹⁸ are independently selected from the group consisting of hydrogen and C₁-C₆ alkyl, or R¹⁷ and R¹⁸ together are a C₂-C₅ alkylene group and with the carbon to which they are attached form a spiro ring of 3 to 6 carbon atoms;
- 15 R¹⁹ is R⁶-phenyl, R⁶-heteroaryl, R⁶-naphthyl, C₂-C₁₀ cycloalkyl, (C₃-C₁₀)cycloalkyl(C₁-C₆alkyl) or (C₁-C₆alkoxy)(C₁-C₆alkyl);
- R²⁰, R²¹ and R²² are independently selected from the group consisting of H and C₁-C₆ alkyl; and
- 20 R²³ is C₁-C₆ alkyl or phenyl.
- 21 The use of claim 16 wherein R is 
- 22 The use of claim 16 wherein R is 
- 23 The use of claim 16 wherein X is -CHOR³, -C(R¹³)(R¹⁹) or -C(=NOR⁴)-.
- 24 The use of claim 18 wherein R³ is pyridyl, R⁴ is (C₁-C₆alkyl), or R¹³ is hydrogen and R¹⁹ is R⁶-phenyl.
- 25 The use of claim 16 wherein R² is R⁷, R⁸, R⁹-phenyl, R⁷, R⁸, R⁹-pyridyl or an N-oxide thereof, or R⁷, R⁸, R⁹-pyrimidinyl.
- 26 The use of claim 20 wherein R² is selected from the group consisting of
-

- 101 -

wherein R⁷ and R⁸ are selected from the group consisting of (C₁-C₆)alkyl, halogen, and -NH₂, and R⁹ is hydrogen.

22. The use of claim 16 for the treatment of Human Immunodeficiency Virus, further comprising one or more antiviral or other agents useful in the treatment of Human Immunodeficiency Virus.

23. The use of claim 22 wherein the antiviral agent is selected from the group consisting of nucleoside reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors and protease inhibitors.

24. The use of claim 16 for the treatment of solid organ transplant rejection, graft v. host disease, inflammatory bowel disease, rheumatoid arthritis or multiple sclerosis, further comprising one or more other agents useful in the treatment of said diseases.

25. A kit comprising in separate containers in a single package pharmaceutical compositions for use in combination to treat Human Immunodeficiency Virus which comprises in one container a pharmaceutical composition comprising an effective amount of a CCR5 antagonist of claim 16 in a pharmaceutically acceptable carrier, and in separate containers, one or more pharmaceutical composition comprising an effective amount of a antiviral or other agent useful in the treatment of Human Immunodeficiency Virus in a pharmaceutically acceptable carrier.

INTERNATIONAL SEARCH REPORT

Int. App. No.
PCT/US 00/1633

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D211/58 C07D417/14 C07D401/14 C07D413/14 A61K31/4468
A61K31/4523 A61P31/12 A61P19/00

B. FIELD(S) SEARCHED

Maximum documentation searched (classification system followed by classification symbols)
IPC 7 C07D A61K A61P

Documentation searched other than maximum documentation to the extent that such documents are included in the field searched

Electronic data bases consulted during the international search (name of data base and, where practical, search terms used)
EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of documents, with indication, where appropriate, of the relevant passages	Relevance to claim No.
X	WO 98 05292 A (SCHERING CORP) 12 February 1998 (1998-02-12) claims 1, 13: examples 6 US 5 889 006 A 30 March 1999 (1999-03-30) cited in the application	1, 2, 25
X	WO 98 06697 A (SCHERING CORP) 19 February 1998 (1998-02-19) claims 1, 11: examples	1, 2, 25
X	WO 98 01425 A (SCHERING CORP) 15 January 1998 (1998-01-15) claims 1, 11: examples	1, 2, 25
	-/-	

* Further documents are listed in the continuation of box C.

* Patent family members are listed in annex.

* Special categories of cited documents:
 "A" document defining the general state of the art which is not considered to be of particular relevance
 "B" earlier document but published on or after the international filing date
 "C" document which may prove decisive on priority claims (i.e. which is cited to establish the publication date of another claim or other special reason (as specified))
 "D" document referring to an oral disclosure, use, exhibition or other means
 "E" document published prior to the international filing date but later than the priority date defined
 "F" document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

1 August 2000

11/08/2000

Name and mailing address of the ISA
 European Patent Office, P.B. 5818 Patenslan 2
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax. (+31-70) 340-2016

Authorized officer
 De Jong, B

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/US 00/11633

C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO 99 04794 A (DATES BRYAN; FINKE PAUL E (US); MACCROSS MALCOLM (US); MERCK & CO I) 1.16</p> <p>February 1999 (1999-02-04)</p> <p>abstract; claim 1</p>	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.
PCT/US 00/11633

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9805292 A	12-02-1998	US 5889006 A	30-03-1999
		AU 3899997 A	25-02-1998
		BR 9711119 A	23-11-1999
		CN 1232462 A	20-10-1999
		CZ 9900366 A	16-06-1999
		EP 0938483 A	01-09-1999
		JP 2000501117 T	02-02-2000
		NO 990551 A	07-04-1999
		PL 331534 A	19-07-1999
		US 6043255 A	28-03-2000
WO 9806697 A	19-02-1998	AU 3973297 A	06-03-1998
		CN 1232453 A	20-10-1999
		CZ 9900433 A	14-07-1999
		EP 0922029 A	16-06-1999
		JP 2000500786 T	25-01-2000
		NO 990671 A	15-04-1999
		PL 331536 A	19-07-1999
WO 9801425 A	15-01-1998	AU 3581097 A	02-02-1998
		CA 2259655 A	15-01-1998
		EP 0912515 A	06-05-1999
		JP 11514671 T	14-12-1999
WO 9904794 A	04-02-1999	AU 8576098 A	16-02-1999
		EP 1003514 A	31-05-2000

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☐ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☒ OTHER: side feet

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.